

Stanford Department of Mathematics Colloquium

NAVIER-STOKES EVOLUTIONS AS SELF-DUAL VARIATIONAL PROBLEMS

Nassif Ghoussoub
UBC

Abstract

We describe how several nonlinear PDEs and evolutions –including stationary and dynamic Navier-Stokes equations– can be formulated and resolved variationally by minimizing energy functionals of the form

$$I(u) = L(u, -\Lambda u) + \langle \Lambda u, u \rangle$$

and

$$I(u) = \int_0^T \left[L(t, u(t), -\dot{u}(t) - \Lambda u(t)) + \langle \Lambda u(t), u(t) \rangle \right] dt + \ell(u(0) - u(T), \frac{u(T) + u(0)}{2})$$

where L is a time-dependent “selfdual Lagrangian” on state space, ℓ is another selfdual “boundary Lagrangian”, and Λ is a nonlinear operator (such as $\Lambda u = \operatorname{div}(u \otimes u)$ in the Navier-Stokes case). However, just like the selfdual Yang-Mills equations, the equations are not obtained via Euler-Lagrange theory, but from the fact that a natural infimum is attained. In dimension 2, we recover the well known solutions for the corresponding initial-value problem as well as periodic and anti-periodic ones, while in dimension 3 we get Leray solutions for the initial-value problems, but also solutions satisfying $u(0) = \alpha u(T)$ for any given α in $(-1, 1)$. It is worth noting that our variational principles translate into Leray’s energy identity in dimension 2 (resp., inequality in dimension 3). Our approach is quite general and does apply to many other situations.

Thursday, October 9.
4:15 p.m.
Bldg. 380, Room 380-W.