

Remarks on mod- $l^n$  Representations,  $l = 3, 5$ 

Brian Conrad

*Department of Mathematics, Harvard University, Cambridge, Massachusetts 02138*E-mail: [bconrad@math.harvard.edu](mailto:bconrad@math.harvard.edu)

and

Siman Wong\*

*Department of Mathematics, Brown University, Providence, Rhode Island 02912*E-mail: [siman@math.brown.edu](mailto:siman@math.brown.edu)*Communicated by K. Rubin*

Received November 9, 1998

Let  $l = 3$  or  $5$ . For any integer  $n > 1$ , we produce an infinite set of triples  $(L, E_1, E_2)$ , where  $L$  is a number field with degree  $l^{3(n-1)}$  over  $\mathbf{Q}$  and  $E_1$  and  $E_2$  are elliptic curves over  $L$  with distinct  $j$ -invariants lying in  $\mathbf{Q}$ , such that the following conditions hold: (1) the pairs of  $j$ -invariants  $\{j(E_1), j(E_2)\}$  are mutually disjoint, (2) the associated mod- $l^n$  representations  $G_L = \text{Gal}(\bar{L}/L) \rightarrow \text{GL}_2(\mathbf{Z}/l^n)$  are surjective, (3) for almost all primes  $\mathfrak{p}$  of  $L$ , we have  $l^n \mid a_{\mathfrak{p}}(E_1)$  if and only if  $l^n \mid a_{\mathfrak{p}}(E_2)$ , and (4) the two representations  $E_i[l^n](\bar{L})$  are not related by twisting by a continuous character  $G_L \rightarrow (\mathbf{Z}/l^n)^\times$ . No such triple satisfying (2)–(4) exists over any number field if we replace  $l$  by a prime larger than 5. The proof depends on determining the automorphisms of the group  $\text{GL}_2(\mathbf{Z}/l^n)$  for  $l = 3, 5$  and analyzing ramification in a branched covering of “twisted” modular curves. © 1999 Academic Press

## 1. INTRODUCTION

Choose a number field  $K$  and fix an algebraic closure  $\bar{K}$  of  $K$ . Denote by  $G_K$  the Galois group  $\text{Gal}(\bar{K}/K)$ . Let  $E_1, E_2$  be elliptic curves over  $K$ ,  $l \in \mathbf{Z}$  a prime,  $n \in \mathbf{Z}$  a positive integer, and fix a basis of  $E_i[l^n](\bar{K})$  over  $\mathbf{Z}/l^n$ . Let

$$\rho_{E_i, l^n}: G_K \rightarrow \text{Aut}(E_i[l^n](\bar{K})) \simeq \text{GL}_2(\mathbf{Z}/l^n)$$

be the resulting mod- $l^n$  representations associated to  $E_i$ , and assume that  $\rho_{E_1, l^n}$  and  $\rho_{E_2, l^n}$  are surjective. Let  $\Sigma$  be a finite set of non-archimedean primes of  $K$  containing all the primes of bad reduction for  $E_1$  and  $E_2$ , as well as all of the primes in  $K$  lying above  $l$ . For any prime  $\mathfrak{p}$  of  $K$  not in  $\Sigma$ , define  $a_{\mathfrak{p}}(E_i)$  to be the trace of the action on the  $l$ -adic Tate module of

\* Current address: Department of Mathematics, University of Massachusetts, Amherst, MA 01003-4515. E-mail: [siman@math.umass.edu](mailto:siman@math.umass.edu).

$E_i$  by an arithmetic Frobenius element at  $\mathfrak{p}$  in  $G_K$ . If  $\rho_{E_1, l^n} \simeq \chi \rho_{E_2, l^n}$  for a continuous character  $\chi: G_K \rightarrow (\mathbf{Z}/l^n)^\times$ , then for all  $\mathfrak{p} \notin \Sigma$ , we have

$$l^n \mid a_{\mathfrak{p}}(E_1) \quad \text{if and only if} \quad l^n \mid a_{\mathfrak{p}}(E_2). \quad (1)$$

By the Chebotarev density theorem, this is equivalent to saying that for all  $g \in G_K$ ,  $\rho_{E_1, l^n}(g)$  has trace 0 if and only if  $\rho_{E_2, l^n}(g)$  has trace 0. It follows from [9, Cor. 1(b)] (and “*Correction to [9]*” below) that if  $l > 5$ , then the condition (1) implies that the  $\rho_{E_i, l^n}$  are equivalent up to twisting by a  $(\mathbf{Z}/l^n)^\times$ -valued continuous character of  $G_K$ . For  $l=3$  or  $l=5$ , and  $n > 1$ , the same conclusion holds for the pair of representations  $G_K \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^{n-1})$  induced from the *surjective*  $\rho_i$  by reduction modulo  $l^{n-1}$ , thanks to [9, Cor. 1(c)]. The proofs depend upon determining the automorphisms of  $\mathrm{PGL}_2(\mathbf{Z}/l^n)$ . For  $l > 5$ , all such automorphisms turn out to be inner, but for  $l=3$  and  $l=5$  there are non-trivial outer automorphisms. In this paper, we exploit these outer automorphisms to produce elliptic curves  $E$  over number fields  $K$  for which the associated mod- $l^n$  representation of  $G_K$  is surjective but is *not* determined (up to twisting) by the set of primes  $\mathfrak{p}$  with  $l^n \mid a_{\mathfrak{p}}(E)$ .

**THEOREM 1.** *Let  $l=3$  or  $5$ , let  $n > 1$ , and let  $K$  be a number field which is linearly disjoint from  $\mathbf{Q}(\zeta_{l^n})$ , where  $\zeta_{l^n}$  is a primitive  $l^n$ th root of unity. There exist infinitely many triples  $(L, E_1, E_2)$  consisting of a finite extension  $L/K$  with degree  $l^{3(n-1)}$  and elliptic curves  $E_1, E_2$  over  $L$  with distinct  $j$ -invariants in  $K$  such that the pairs  $\{j(E_1), j(E_2)\}$  are mutually disjoint, the corresponding mod- $l^n$  representations  $\rho_{E_1, l^n}, \rho_{E_2, l^n}: G_L \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^n)$  satisfy the condition (1) and are surjective, and  $\rho_{E_1, l^n}$  and  $\rho_{E_2, l^n}$  are not equivalent up to twisting by any continuous character  $G_L \rightarrow (\mathbf{Z}/l^n)^\times$ . In fact, infinitely many such triples  $\tau = (L, E_1, E_2)$  can be chosen so that each pair of representations  $\rho_{E_1, l^n}$  and  $\rho_{E_2, l^n}$  has the same common splitting field  $L_\tau$  over  $L$  and as we vary  $\tau$ , no prime of  $K$  away from  $l$  with norm  $> (l^2 - 3)/2$  is ramified in more than one of the  $L_\tau$ 's.*

In view of our remarks above, for any triple  $(L, E_1, E_2)$  in the theorem, the mod- $l^{n+1}$  representations  $G_L \rightarrow \mathrm{GL}_2(\mathbf{Z}/l^{n+1})$  arising from  $E_1$  and  $E_2$  cannot *both* be surjective. To prove the theorem, we use a non-trivial outer automorphism of  $\mathrm{PGL}_2(\mathbf{Z}/l^n)$  in order to construct a non-trivial determinant-preserving outer automorphism  $\varphi$  of  $\mathrm{GL}_2(\mathbf{Z}/l^n)$  which takes trace zero matrices to trace zero matrices. If  $\rho$  is a *surjective* mod- $l^n$  representation of an elliptic curve  $E$  over a number field  $K$ , then  $\rho$  and  $\rho' = \varphi \circ \rho$  have cyclotomic determinant and are not equivalent up to twists. Moreover, for all but finitely many primes  $\mathfrak{p}$  of  $K$ ,  $\rho$  and  $\rho'$  are unramified at  $\mathfrak{p}$  and  $l^n \mid \mathrm{trace}(\rho(\mathrm{Frob}_{\mathfrak{p}}))$  if and only if  $l^n \mid \mathrm{trace}(\rho'(\mathrm{Frob}_{\mathfrak{p}}))$ , where  $\mathrm{Frob}_{\mathfrak{p}}$  is an

arithmetic Frobenius element at  $\mathfrak{p}$  in  $G_K$ . We want to realize  $\rho'$  as the mod- $l^n$  representation of an elliptic curve  $E'$  over  $K$ . This step will require enlarging  $K$  a small amount to an extension  $L$ , but we will be able to slightly control ramification in  $L/K$ .

Here is how we will find  $E'$ . There is a proper smooth curve  $X(\rho')$  over  $K$  which, roughly speaking, classifies elliptic curves whose mod- $l^n$  representation is isomorphic to  $\rho'$ . In particular, over  $\bar{K}$  there is an isomorphism

$$X(\rho') \times_K \bar{K} \simeq X(l^n) \times_{\mathbf{Z}[1/l]} \bar{K},$$

where  $X(l^n)$  denotes the compactified full level  $l^n$  moduli scheme over  $\mathbf{Z}[1/l]$  in the sense of [5, Sects. 8.6ff.], so  $X(\rho')$  is *not* geometrically connected over  $K$ . However, since the determinant of  $\rho'$  is cyclotomic, the connected components of  $X(\rho')$  are geometrically connected over  $K$ . Let  $\bar{\rho}'$  be the mod- $l$  reduction of  $\rho'$ . "Reduction mod  $l$ " on Galois representations induces a finite flat map  $X(\rho') \rightarrow X(\bar{\rho}')$  over  $K$  whose base change to  $\bar{K}$  is the usual projection  $X(l^n) \times_{\mathbf{Z}[1/l]} \bar{K} \rightarrow X(l) \times_{\mathbf{Z}[1/l]} \bar{K}$ .

For  $l=3$  and  $5$ , an argument of Mazur shows that the connected components of  $X(\bar{\rho}')$  have rational points and so are non-canonically isomorphic to  $\mathbf{P}_K^1$ . Thus, we can regard the connected components of  $X(\rho')$  as branched covers of  $\mathbf{P}_K^1$  which are geometrically connected over  $K$ . We find the desired elliptic curves in Theorem 1 by looking in the fibers on  $X(\rho')$  over well-chosen  $K$ -rational points on the connected components  $\mathbf{P}_K^1$  of  $X(\bar{\rho}')$ . We do not know if it is sufficient to only look at  $K$ -rational points on  $X(\rho')$  (of which there are only finitely many, by Faltings' Theorem), and this is why we cannot precisely control the number fields over which our examples occur.

*Correction to [9].* S. W. would like to take this opportunity to correct a confusing terminology mistake in [9], which is needed in the present paper. Let  $\mathcal{O}$  be a complete local ring with maximal ideal  $\lambda$ . Consider two continuous representations  $\rho_1, \rho_2: G_K \rightarrow \mathrm{GL}_n(\mathcal{O})$  which are unramified outside of a finite set of places  $\Sigma$  of  $K$ . For any  $\mathfrak{p} \notin \Sigma$ , define  $a_i(\mathfrak{p}) = \mathrm{trace} \rho_i(\mathrm{Frob}_{\mathfrak{p}})$ . In [9, Sect. 1] (see in particular the displayed equation (1) there),  $\rho_1$  and  $\rho_2$  are defined to be " $\lambda$ -adically close at the supersingular primes" if there is a positive integer  $N_0$  such that whenever *both*  $a_i(\mathfrak{p})$  lie in  $\lambda^{N_0}$ , one has for all  $w \geq N_0$  that  $a_1(\mathfrak{p}) \in \lambda^w$  if and only if  $a_2(\mathfrak{p}) \in \lambda^w$ . This definition is inadequate for the proofs in [9], and is automatically satisfied whenever  $\lambda^{N_0} = 0$  (a case of interest for the present paper)! The definition of  $\lambda$ -adic closeness should have been modified to require that if *one of the two*  $a_i(\mathfrak{p}) \in \lambda^{N_0}$ , then for any  $w \geq N_0$ ,  $a_1(\mathfrak{p}) \in \lambda^w$  if and only if  $a_2(\mathfrak{p}) \in \lambda^w$ . Note, for example, that this is a non-trivial condition even if  $\lambda^{N_0} = 0$ .

It is only under this modified definition of  $\lambda$ -adic closeness that the arguments in [9] yield the results as claimed there. However, the statement

of [9, Lemma 7] needs to be slightly modified. Beginning with the phrase *Suppose one of the following holds...*, the lemma should be replaced by the following:

Suppose one of the following holds:

- $n$  is even and either  $k \neq \mathbf{F}_2$  or 2 is not a zero-divisor in  $\mathcal{O}$ ; or
- $n \geq 5$  is odd and either  $k \neq \mathbf{F}_3$  or 3 is not a zero-divisor in  $\mathcal{O}$ ; or
- $n = 3$  and  $k \neq \mathbf{F}_2$ ,  $k \neq \mathbf{F}_3$ .

Then there exists an automorphism  $\varphi$  of  $\mathrm{PGL}_n(\mathcal{O})$  such that  $\varphi \circ \tilde{\rho}_2 = \tilde{\rho}_1$ .

Suppose instead that  $n$  is even and  $k = \mathbf{F}_2$ , or that  $n = 3$  and  $k = \mathbf{F}_3$ . Let  $p$  denote the characteristic of  $k$  and let  $\mathfrak{a}$  denote the annihilator of  $p$  in  $\mathcal{O}$ . Then the analogous conclusion holds for the pair of representations  $G_K \rightarrow \mathrm{PGL}_n(\mathcal{O}/\mathfrak{a})$  induced from the  $\tilde{\rho}_i$ .

## 2. BRANCHED COVERS OF $\mathbf{P}_K^1$

In this section, we recall some results related to the Hilbert Irreducibility Theorem, stated in a geometric form.

Let  $K$  be a number field and let  $\pi: X \rightarrow \mathbf{P}_K^1$  be a finite map, where  $X$  is a smooth connected curve over  $K$ . The Hilbert Irreducibility Theorem says that for infinitely many  $K$ -rational points  $a \in \mathbf{P}_K^1$ , the fiber  $\pi^{-1}(a)$  has the form  $\pi^{-1}(a) \simeq \mathrm{Spec}(L_a)$  for a finite extension field  $L_a/K$ . In more algebraic terms, if we identify  $K(\mathbf{P}_K^1) \simeq K(t)$  and we choose a primitive element for the finite separable extension  $K(X)/K(\mathbf{P}_K^1)$  of function fields, then  $K(X) \simeq K(t)[Y]/(f)$  for some monic  $f \in K(t)[Y]$ . The Hilbert Irreducibility Theorem in the geometric form just given is equivalent to the statement that for infinitely many  $t_0 \in K$ , the polynomial  $f(t_0, Y) \in K[Y]$  is irreducible, in which case  $L_{t_0} = K[Y]/(f(t_0, Y))$ . Of course, we avoid the finitely many  $t_0 \in K$  where some coefficient of  $f$  in  $K(t)$  has a pole.

We will need a milder stronger formulation, which is well-known:

**LEMMA 1.** *Let  $\pi$  be as above and choose a finite extension  $E/K$ . Assume that  $X$  is geometrically connected over  $K$ , or more generally that  $E$  is linearly disjoint (over  $K$ ) from the algebraic closure of  $K$  in  $K(X)$ . Then there exist infinitely many  $K$ -rational points  $a \in \mathbf{P}_K^1$  for which  $\pi^{-1}(a) \simeq \mathrm{Spec}(L_a)$  for a finite extension  $L_a/K$  which is linearly disjoint from  $E$  over  $K$ . In other words,  $\pi^{-1}(a) \times_K E$  is irreducible for infinitely many  $K$ -rational points  $a \in \mathbf{P}_K^1$ .*

*Proof.* Since  $E/K$  is a finite separable extension, by [6, Prop 3.3, Sect. 9] every Hilbert set in  $E$  contains a Hilbert set in  $K$ . Put in more algebraic terms, for any irreducible monic polynomial  $f \in E(t)[Y]$ , there exists an irreducible

monic polynomial  $g_f \in K(t)[Y]$  such that for all but finitely many  $t_0 \in K$ ,  $f(t_0, Y) \in E[Y]$  is irreducible whenever  $g_f(t_0, Y) \in K[Y]$  is irreducible. Thus, by the Hilbert Irreducibility Theorem for the number field  $K$  and the polynomial  $g_f \in K(t)[Y]$ , we conclude that for any irreducible monic  $f \in E(t)[Y]$ , there are infinitely many  $t_0 \in K$  (rather than just  $t_0 \in E$ ) such that  $f(t_0, Y) \in E[Y]$  is irreducible. In particular, for any irreducible monic  $f \in K(t)[Y]$  which remains irreducible in  $E(t)[Y]$ , there are infinitely many  $t_0 \in K$  so that  $f(t_0, Y)$  is irreducible in  $E[Y]$ . Of course, this is just the usual proof that a finite (separable) extension of a Hilbertian field is again Hilbertian.

In order to use this to deduce the lemma, we just have to show that if we choose an isomorphism  $K(X) \simeq K(t)[Y]/(f)$  for some irreducible monic  $f \in K(t)[Y]$ , then  $f$  is irreducible in  $E(t)[Y]$ . It is not difficult to show that this is equivalent to the irreducibility of  $X \times_K E$ , or even the connectedness of  $X \times_K E$  (by smoothness). If  $K'$  denotes the algebraic closure of  $K$  in  $K(X)$  then  $X$  is naturally a proper smooth curve over  $K'$  and is geometrically connected as such [3, IV<sub>2</sub>, 4.5.15]. Since  $X \times_K E = X \times_{K'} \text{Spec}(K' \otimes_K E)$  and  $K' \otimes_K E$  is a field by the linear disjointness hypothesis, it follows that  $X \times_K E$  is connected. ■

### 3. AUTOMORPHISMS OF $\text{GL}_2(\mathbf{Z}/l^n)$

**LEMMA 2.** *Let  $R$  be a local ring with residue field  $k$  and maximal ideal  $\mathfrak{m}$ . The natural map  $\text{SL}_n(R) \rightarrow \text{SL}_n(k)$  is surjective. The same holds with  $\text{PSL}_n$  replaced by  $\text{PSL}_n$ ,  $\text{PGL}_n$  and  $\text{GL}_n$ .*

*Proof.* Given a matrix  $A = (a_{ij})$  in  $\text{SL}_n(k)$ , let  $\mathfrak{a} = (\alpha_{ij})$  be an  $n \times n$  matrix over  $R$  with  $\alpha_{ij} \bmod \mathfrak{m} = a_{ij}$  for all  $i, j$ . Denote by  $\alpha_{ij}$  the  $(n-1) \times (n-1)$  matrix obtained by removing the  $i$ th row and the  $j$ th column of  $\mathfrak{A}$ . Define  $A_{ij}$  similarly. Then

$$\sum_{j=1}^n (-1)^j \alpha_{1j} \det(\mathfrak{a}_{1j}) = \det(\mathfrak{a}) \equiv 1 \pmod{\mathfrak{m}}. \quad (2)$$

If we fix the entries  $\alpha_{ij}$  with  $i \geq 2$ , then any lift  $\mathfrak{a}$  of  $A$  with these  $\alpha_{ij}$  for  $i > 2$  gives rise to a solution mod  $\mathfrak{m}$  of the linear equation (2). Moreover, since  $\det(\mathfrak{a}_{1j}) \bmod \mathfrak{m} = \det(A_{1j})$  for all  $j$ , at least one of the  $\det(\mathfrak{a}_{ij})$  is a unit. Thus, we can easily find elements  $\alpha_{11}, \dots, \alpha_{1n}$  in  $R$  so that the left side of (2) is equal to 1 in  $R$ . This takes care of the lemma for  $\text{SL}_n$ ; the other cases are similar. ■

**LEMMA 3.** *Let  $R$  be a Noetherian local ring with maximal ideal  $\mathfrak{m}$  and finite residue field  $k$  with characteristic  $l > 0$ . Denote by  $K_n$  and  $L_n$  the kernel*

of the natural maps from  $\mathrm{PSL}_n(R)$  to  $\mathrm{PSL}_n(R/\mathfrak{m})$  and  $\mathrm{PSL}_n(R/\mathfrak{m}^2)$ , respectively. Let  $M \subset \mathrm{PSL}_n(R)$  be a normal subgroup such that  $ML_n/L_n$  is a finite  $l$ -group. Then  $M \subset K_n$ .

The same conclusion holds if  $l \nmid n$  and if we replace  $\mathrm{PSL}_n$  by  $\mathrm{PGL}_n$ .

*Remark 1.* The group  $ML_n/L_n$  is always finite: it is a subgroup of  $\mathrm{PSL}_n(R)/K_n$ , which injects into  $\mathrm{PSL}_n(R/\mathfrak{m}^2)$ , which is finite since  $k$  is finite and  $\mathfrak{m}$  is finitely generated.

*Proof.* We first deal with the case of  $\mathrm{PSL}_n$ . Then there are no non-trivial normal  $l$ -subgroups in  $\mathrm{PSL}_n(k)$ : for  $n \neq 2$  or  $k \neq \mathbf{F}_2, \mathbf{F}_3$  this follows from the simplicity of  $\mathrm{PSL}_n(k)$ , and the remaining cases follow from the isomorphisms  $\mathrm{PSL}_2(\mathbf{F}_2) \simeq S_3$  and  $\mathrm{PSL}_2(\mathbf{F}_3) \simeq A_4$ .

Since  $\mathfrak{m}/\mathfrak{m}^2$  is a finite-dimensional  $k$ -vector space,  $K_n/L_n$  is a finite elementary  $l$ -group, and hence so is  $MK_n/ML_n$ . The exact sequence

$$1 \rightarrow ML_n/L_n \rightarrow MK_n/L_n \rightarrow MK_n/ML_n \rightarrow 1$$

and the hypothesis on  $M$  then imply that  $MK_n/L_n$ , and hence  $MK_n/K_n$ , is a finite  $l$ -group. The latter is a normal  $l$ -subgroup of  $\mathrm{PSL}_n(R)/K_n$ , which by Lemma 2 is isomorphic to  $\mathrm{PSL}_n(k)$ . Thus  $MK_n = K_n$ , as desired.

The quotient group  $\mathrm{PGL}_n(R)/\mathrm{PSL}_n(R) \simeq R^\times/R^{\times n}$  has exponent dividing  $n$ , so the above argument applies to  $\mathrm{PGL}_n$  if  $l \nmid n$ .

**COROLLARY 1.** *Let  $R$  be a Noetherian local ring with maximal ideal  $\mathfrak{m}$  and a finite residue field  $k$  with characteristic  $l > 0$ . Define  $K_n$  as in Lemma 3. Every automorphism  $\varphi$  of  $\mathrm{PSL}_n(R)$  (resp.  $\mathrm{PGL}_n(R)$  with  $l \nmid n$ ) takes  $K_n$  to itself, thereby giving an automorphism  $\bar{\varphi}$  of  $\mathrm{PSL}_n(k)$  (resp.  $\mathrm{PGL}_n(k)$ ) such that  $\bar{\varphi}(\bar{g}) = \overline{\varphi(g)}$  for all  $g \in \mathrm{PSL}_n(R)$  (resp.  $g \in \mathrm{PGL}_n(R)$ ), where  $\overline{(\cdot)}$  denotes the image under the natural map  $\mathrm{PSL}_n(R) \rightarrow \mathrm{PSL}_n(k)$  (resp.  $\mathrm{PGL}_n(R) \rightarrow \mathrm{PGL}_n(k)$ ).*

*Proof.* Apply Lemma 3 to  $M = \varphi(K_n)$ . ■

For the rest of this section, fix a prime  $l$ , let  $\alpha \in (\mathbf{Z}/l^n)^\times$  be a choice of generator of the unique cyclic subgroup order  $l-1$ , and let  $\Gamma$  be the subgroup of  $GL_2(\mathbf{Z}/l^n)$  generated by  $\begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}$  and  $SL_2(\mathbf{Z}/l^n)$ . Thus,  $\Gamma$  is abstractly a semi-direct product  $\mathbf{Z}/(l-1) \rtimes SSL_2(\mathbf{Z}/l^n)$ , where the  $\mathbf{Z}/(l-1)$  is generated by  $\begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}$ . Since  $SL_2(\mathbf{Z}/l^n)$  contains all elements in  $\Gamma$  with  $l$ -power order and it is generated by such elements (e.g.,  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ ), we see that  $SL_2(\mathbf{Z}/l^n)$  is stable under  $\mathrm{Aut}(\Gamma)$ . The natural map  $\Gamma \rightarrow GL_2(\mathbf{Z}/l)$  is clearly surjective, and if  $l > 2$ , then the scalar matrices in  $\Gamma$  are those of order dividing  $l-1$ . Also, note that if  $l > 2$ , then the restriction of the canonical map  $GL_2(\mathbf{Z}/l^n) \xrightarrow{\pi} \mathrm{PGL}_2(\mathbf{Z}/l^n)$  to  $\Gamma$  is surjective.

LEMMA 4. *If  $l > 2$ , then every automorphism of  $\mathrm{PGL}_2(\mathbf{Z}/l^n)$  lifts to an automorphism of  $\Gamma$ .*

*Proof.* Choose an automorphism  $\varphi$  of  $\mathrm{PGL}_2(\mathbf{Z}/l^n)$ . Since

$$\begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix} = \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}^2 \begin{pmatrix} \alpha^{-1} & 0 \\ 0 & \alpha \end{pmatrix},$$

we see that  $H = \ker(\pi|_{\Gamma})$  is a cyclic group (of scalar matrices) of order  $l-1$ , and

$$1 \rightarrow H \rightarrow \Gamma \xrightarrow{\pi} \mathrm{PGL}_2(\mathbf{Z}/l^n) \rightarrow 1 \quad (3)$$

is a central extension, corresponding to a cohomology class  $\phi \in H^2(\mathrm{PGL}_2(\mathbf{Z}/l^n), H)$  (the surjectivity of  $\pi$  in (3) requires  $l > 2$ ). Since the automorphism group of the cyclic group  $H$  is commutative, an easy calculation shows that the automorphism  $\varphi$  of  $\mathrm{PGL}_2(\mathbf{Z}/l^n)$  lifts to an automorphism of  $\Gamma$  if and only if  $\varphi^*(\phi) \in H^2(\mathrm{PGL}_2(\mathbf{Z}/l^n), H)$  is equal to the image  $\varphi_H(\phi)$  for some automorphism  $\varphi_H: H \simeq H$ . The point is that when such a  $\varphi_H$  exists, there is a lift  $\tilde{\varphi}$  of  $\varphi$  to an endomorphism of  $\Gamma$  which induces the automorphism  $\varphi_H$  on  $H$ . A simple diagram chase then shows that  $\tilde{\varphi}$  is actually an automorphism of  $\Gamma$ .

The only possibilities for  $\varphi_H$  are multiplication by  $m \in (\mathbf{Z}/(l-1))^\times$ , and if  $\varphi^*(\phi) = m\phi$  for some  $m \in \mathbf{Z}/(l-1)$ , then (by the Chinese Remainder Theorem)  $m$  can be chosen to lie in  $(\mathbf{Z}/(l-1))^\times$  (since  $\varphi^*$  is an automorphism). Thus,  $\varphi$  lifts to an automorphism of  $\Gamma$  if and only if the cohomology class  $\varphi^*(\phi) = \varphi_H(\phi)$  for some endomorphism  $\varphi_H$  of the group  $H$ . By an argument in terms of central extensions, it is clear that the elements of the form  $\varphi_H(\phi)$  for variable  $\varphi_H$  are precisely the elements in the kernel of  $\pi^*: H^2(\mathrm{PSL}_2(\mathbf{Z}/l^n), H) \rightarrow H^2(\Gamma, H)$ . Thus,  $\varphi$  lifts to an automorphism of  $\Gamma$  if and only if  $(\varphi \circ \pi)^* \phi = \pi^* \varphi^* \phi = 0$  in  $H^2(\Gamma, H)$ . We will show that  $(\varphi \circ \pi)^* \phi = 0$ .

Let  $K = \ker(\Gamma \rightarrow \mathrm{GL}_2(\mathbf{Z}/l))$  and let  $P = \ker(\Gamma \rightarrow \mathrm{PGL}_2(\mathbf{Z}/l))$ . Since (3) is a central extension,  $P$  and  $K$  act trivially on  $H$ . Also, since  $K$  is a finite  $l$ -group and  $H$  has order prime to  $l$ ,  $H^i(K, H) = 0$  for all  $i > 0$ . Since  $\pi(P)$  is the kernel of the natural map  $\mathrm{PGL}_2(\mathbf{Z}/l^n) \rightarrow \mathrm{PGL}_2(\mathbf{Z}/l)$ , it follows from Lemma 3 that  $\varphi$  takes  $\pi(P)$  isomorphically back to itself. The induced automorphism  $\bar{\varphi}$  of  $\mathrm{PGL}_2(\mathbf{Z}/l^n)/\pi(P) \simeq \mathrm{PGL}_2(\mathbf{Z}/l)$  is exactly the map in Corollary 1, so composing the map  $\Gamma/K \rightarrow \mathrm{PGL}_2(\mathbf{Z}/l^n)/\pi(K)$  (induced by  $\pi$ ) with the projection  $\mathrm{PGL}_2(\mathbf{Z}/l^n)/\pi(K) \rightarrow \mathrm{PGL}_2(\mathbf{Z}/l^n)/\pi(P)$  and the automorphism  $\bar{\varphi}$ , we get a map of groups  $\psi: \Gamma/K \rightarrow \mathrm{PGL}_2(\mathbf{Z}/l^n)/\pi(P)$ . Using the identification  $\Gamma/K \simeq \mathrm{GL}_2(\mathbf{Z}/l)$ , this map  $\psi$  is exactly the composite

of the canonical projection  $\bar{\pi}: GL_2(\mathbf{Z}/l) \rightarrow PGL_2(\mathbf{Z}/l)$  and the automorphism  $\bar{\varphi}$  of  $PGL_2(\mathbf{Z}/l)$ . The kernel of  $\bar{\pi}$  is just the mod  $l$  “reduction” of  $H$ , which is canonically identified with  $H$ , due to how  $H$  is defined.

Functoriality and the inflation-restriction sequence therefore yield the commutative diagram

$$\begin{CD}
 H^2(PGL_2(\mathbf{Z}/l^n)/\pi(P), H) @>\beta>> H^2(PGL_2(\mathbf{Z}/l^n), H) \\
 @VV\psi^*V @VV(\varphi \circ \pi)^*V \\
 H^2(\Gamma/K, H) @>\sim>> H^2(\Gamma, H).
 \end{CD} \tag{4}$$

in which the bottom row is an isomorphism and the left column is identified with the map

$$(\bar{\varphi} \circ \bar{\pi})^*: H^2(PGL_2(\mathbf{Z}/l), H) \simeq H^2(GL_2(\mathbf{Z}/l), H).$$

The cohomology class  $\bar{\phi}$  in  $H^2(PGL_2(\mathbf{Z}/l), H)$  corresponding to the central extension

$$1 \rightarrow H \rightarrow GL_2(\mathbf{Z}/l) \xrightarrow{\bar{\pi}} PGL_2(\mathbf{Z}/l) \rightarrow 1 \tag{5}$$

satisfies  $\beta(\bar{\phi}) = \phi$ . Thus,  $(\varphi \circ \pi)^* \phi = 0$  if and only if  $(\bar{\varphi} \circ \bar{\pi})^* (\bar{\phi}) = 0$ , which is to say that the automorphism  $\bar{\varphi}$  of  $PGL_2(\mathbf{Z}/l)$  can be lifted to an automorphism of  $GL_2(\mathbf{Z}/l)$ . The liftability of all such automorphisms is classical [2, Thm. V.5]. ■

For any ring  $R$ , if  $\varphi$  is an automorphism of  $GL_2(R)$ , then  $\varphi$  takes the diagonal matrices of  $GL_2(R)$  to themselves (since these matrices constitute the center of  $GL_2(R)$ ). Thus  $\varphi$  induces a group homomorphism  $r_\varphi: R^\times \rightarrow R^\times$ .

**LEMMA 5.** *Let  $R$  be a local ring whose residue field is not  $\mathbf{F}_2$ . Then every automorphism  $\varphi$  of  $GL_2(R)$  takes  $SL_2(R)$  to itself. Moreover, if  $\varphi_1$  and  $\varphi_2$  are two automorphisms of  $GL_2(R)$ , then  $\varphi_1$  and  $\varphi_2$  coincide on  $SL_2(R)$  if and only if there is a map of groups  $\lambda: R^\times \rightarrow R^\times$  such that  $\varphi_1(g) = \lambda(\det(g)) \varphi_2(g)$  for all  $g \in GL_2(R)$ . Conversely, for any map of groups  $\lambda: R^\times \rightarrow R^\times$  and any automorphism  $\varphi$  of  $GL_2(R)$ ,  $\lambda^2 r_\varphi$  is an automorphism of  $R^\times$  if and only if the map  $g \mapsto \lambda(\det(g)) \varphi(g)$  defines an automorphism of  $GL_2(R)$ .*

*Proof.* For a local ring  $R$  as above, the commutator subgroup of  $GL_2(R)$  is  $SL_2(R)$  [1, Thm 4.1, Prop 9.2]. The first part of the lemma then follows, and any group map  $GL_2(R) \rightarrow R^\times$  must factor through the determinant map. To prove the second part, it suffices to consider an endomorphism  $\varphi$  of the group  $GL_2(R)$  such that  $\varphi$  is the identity on  $SL_2(R)$ , and to show that  $\varphi(g) = \lambda(\det(g))g$  for all  $g \in GL_2(R)$ , where



$\lambda: R^\times \rightarrow R^\times$  is some map of groups. Pick an element  $\mu \in R^\times$  and write  $\varphi\left(\begin{smallmatrix} 1 & 0 \\ 0 & \mu \end{smallmatrix}\right) = \begin{pmatrix} x & y \\ z & w \end{pmatrix}$ . We have the identities

$$\begin{aligned} \begin{pmatrix} 1 & 0 \\ 0 & \mu \end{pmatrix} \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \mu \end{pmatrix}^{-1} &= \begin{pmatrix} 1 & \lambda/\mu \\ 0 & 1 \end{pmatrix}, \\ \begin{pmatrix} 1 & 0 \\ 0 & \mu \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \lambda & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \mu \end{pmatrix}^{-1} &= \begin{pmatrix} 1 & 0 \\ \lambda\mu & 1 \end{pmatrix}. \end{aligned}$$

Since  $\varphi$  is trivial on  $\mathrm{SL}_2(R)$ , applying  $\varphi$  to these identities and comparing the entries yields  $y = z = 0$  and  $\mu = w/x$ . Thus  $\varphi\left(\begin{smallmatrix} 1 & 0 \\ 0 & \mu \end{smallmatrix}\right) = \lambda(\mu)\left(\begin{smallmatrix} 1 & 0 \\ 0 & \mu \end{smallmatrix}\right)$  for some  $\lambda(\mu) \in R^\times$ . Since  $\varphi$  is multiplicative,  $\lambda$  is an endomorphism of the group  $R^\times$ . Every element  $g$  of  $\mathrm{GL}_2(R)$  can be written uniquely as  $g'\left(\begin{smallmatrix} 1 & 0 \\ 0 & \det(g) \end{smallmatrix}\right)$  with  $g' \in \mathrm{SL}_2(R)$ , so  $\varphi(g) = \lambda(\det(g))g$  for all  $g \in \mathrm{GL}_2(R)$ .

Finally, let  $\varphi$  be an automorphism of the group  $\mathrm{GL}_2(R)$  and let  $\lambda: R^\times \rightarrow R^\times$  be a map of groups. Then  $\varphi_\lambda: g \mapsto \lambda(\det(g))\varphi(g)$  is an endomorphism of  $\mathrm{GL}_2(R)$  which induces an automorphism on  $\mathrm{SL}_2(R)$ . Suppose

$$\varphi_\lambda(g) = \varphi_\lambda(h) \tag{6}$$

for some  $g, h \in \mathrm{GL}_2(R)$ . Then  $\lambda(\det(g))\lambda(\det(h))^{-1}\left(\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix}\right) = \varphi(g^{-1}h)$ . Since  $\varphi$  induces an automorphism of the scalar matrices, we have  $h = g\left(\begin{smallmatrix} s & 0 \\ 0 & s \end{smallmatrix}\right)$  for some  $s \in R^\times$ . Since  $\varphi_\lambda$  is a homomorphism, it follows from (6) that  $\varphi_\lambda(h) = \varphi_\lambda(g)\varphi_\lambda\left(\begin{smallmatrix} s & 0 \\ 0 & s \end{smallmatrix}\right)$ , and hence  $\lambda^2(s)r_\varphi(s) = 1$ . Conversely, if  $(\lambda^2 r_\varphi)(s) = 1$  for some  $s \in R^\times$ , then  $\varphi_\lambda\left(\begin{smallmatrix} s & 0 \\ 0 & s \end{smallmatrix}\right) = 1$ . Thus  $\varphi_\lambda$  is injective if and only if  $\lambda^2 r_\varphi$  is injective.

Denote by  $\mathcal{S}$  the subgroup of  $\mathrm{GL}_2(R)$  generated by  $\mathrm{SL}_2(R)$  and by the scalar matrices. Note that  $\varphi_\lambda$  takes  $\mathcal{S}$  to itself, and induces an automorphism of  $\mathcal{S}$  if  $\varphi_\lambda$  is an automorphism. Since  $\varphi_\lambda\left(\begin{smallmatrix} \beta & 0 \\ 0 & \beta \end{smallmatrix}\right) = \lambda^2 r_\varphi(\beta)\left(\begin{smallmatrix} 1 & 0 \\ 0 & 1 \end{smallmatrix}\right)$ , we conclude that  $\lambda^2 r_\varphi$  is an automorphism (of the scalar matrices) if and only if  $\varphi_\lambda$  induces an automorphism of  $\mathcal{S}$ . Thus  $\varphi_\lambda$  always induces a map  $\tilde{\varphi}_\lambda$  on  $\mathrm{GL}_2(R)/\mathcal{S}$ , and  $\varphi_\lambda$  is an automorphism if and only if  $\lambda^2 r_\varphi$  is an automorphism and  $\tilde{\varphi}_\lambda$  is surjective on  $\mathrm{GL}_2(R)/\mathcal{S}$ . But the action of  $\tilde{\varphi}_\lambda$  on  $\mathrm{GL}_2(R)/\mathcal{S}$  is the same as that of  $\varphi$  on  $\mathrm{GL}_2(R)/\mathcal{S}$ , which is surjective since  $\varphi$  is an automorphism of  $\mathrm{GL}_2(R)$ , so we are done. ■

**LEMMA 6.** *Let  $l = 3$  or  $5$ , and let  $n > 1$ . Let  $v, t \in \mathbf{Z}/l^n$  be divisible by  $l^{n-1}$ , with  $t = 0$  or  $3$  if  $l = 3$  and  $n = 2$ . Then the following*

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} &\mapsto \begin{pmatrix} 1 & 1 \\ t & 1+t \end{pmatrix}, & \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} &\mapsto \begin{pmatrix} 0 & t-1 \\ t+1 & 0 \end{pmatrix}, \\ \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} &\mapsto \begin{pmatrix} \alpha & v \\ v & 1 \end{pmatrix}, & \begin{pmatrix} \beta & 0 \\ 0 & \beta \end{pmatrix} &\mapsto \begin{pmatrix} \beta & 0 \\ 0 & \beta \end{pmatrix} \end{aligned}$$

determine a unique automorphism  $\phi_{v,t}$  of  $\text{GL}_2(\mathbf{Z}/l^n)$ , and  $\phi_{v,t}$  is determinant-preserving. When  $v \neq 0$  or  $t \neq 0$ , then  $\phi_{v,t}$  is not an inner automorphism.

Every automorphism of  $\text{GL}_2(\mathbf{Z}/l^n)$  has the form

$$\phi_{v,t,\lambda,h}: g \mapsto \lambda(\det(g)) h\phi_{v,t}(g) h^{-1}$$

for  $h \in \text{GL}_2(\mathbf{Z}/l^n)$  and a map of groups  $\lambda: (\mathbf{Z}/l^n)^\times \rightarrow (\mathbf{Z}/l^n)^\times$ . Such automorphisms take elements with trace zero to elements with trace zero. Finally, for any  $v, t, \lambda, h$  as above, the map  $\phi_{v,t,\lambda,h}$  is an automorphism of  $\text{GL}_2(\mathbf{Z}/l^n)$  if and only if  $\lambda^2(a) \neq a^{-1}$  for all  $a \in (\mathbf{Z}/l^n)^\times$  with  $a \neq 1$ .

*Proof.* With  $t$  and  $v$  as in the lemma, it follows from [9, Thm. 3] and our hypothesis that  $l = 3$  or  $l = 5$  that

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & 1 \\ t & t+1 \end{pmatrix}, \quad \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \mapsto \begin{pmatrix} 0 & t-1 \\ t+1 & 0 \end{pmatrix}, \quad \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} \alpha & v \\ v & 1 \end{pmatrix} \tag{7}$$

determines a unique automorphism  $\varphi_{v,t}$  of  $\text{PGL}_2(\mathbf{Z}/l^n)$ , and that every automorphism of  $\text{PGL}_2(\mathbf{Z}/l^n)$  is the compositum of an inner one with some  $\varphi_{v,t}$ . Moreover, by [9, Cor 2] and our hypothesis that  $l = 3$  or  $l = 5$ , the first two conditions in (7) determine a unique automorphism of  $\text{SL}_2(\mathbf{Z}/l^n)$ . Since  $-1 \in (\mathbf{Z}/l^n)^\times$  does not have  $l$ -power order, by Lemma 4 and our earlier observation that  $\text{SL}_2(\mathbf{Z}/l^n) \subseteq \Gamma$  is stable under  $\text{Aut}(\Gamma)$  we see that there exists an automorphism  $\Phi_{v,t}$  of  $\Gamma$  satisfying the first two conditions of (7), with

$$\Phi_{v,t} \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \alpha & v \\ v & 1 \end{pmatrix} \begin{pmatrix} \gamma & 0 \\ 0 & \gamma \end{pmatrix}$$

for some  $\gamma \in (\mathbf{Z}/l^n)^\times$ . Since  $\begin{pmatrix} \alpha & v \\ v & 1 \end{pmatrix}$  has order  $l-1$  in  $\text{GL}_2(\mathbf{Z}/l^n)$ , we have  $\gamma^{l-1} = 1$ , so we can write

$$\Phi_{v,t} \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} = \det \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix}^A \begin{pmatrix} \alpha & v \\ v & 1 \end{pmatrix}$$

for some  $A \in \mathbf{Z}$ . The scalars in  $\Gamma$  are the powers of  $\begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}$  since  $l > 2$ , and it is easy to compute that

$$\begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix} = \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^{-1},$$

so  $\Phi_{v,t}$  acts as multiplication by  $\alpha^{2A}$  on  $\begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}$ . Since  $\text{GL}_2(\mathbf{Z}/l^n)$  is generated by the commuting subgroups  $\Gamma$  and  $(\mathbf{Z}/l^n)^\times$  (i.e., the scalar matrices), we can extend  $\det^{-A} \Phi_{v,t}$  to an endomorphism  $\phi_{v,t}$  of the group  $\text{GL}_2(\mathbf{Z}/l^n)$  by

letting it acts trivially on the scalar matrices. It is easy to see that  $\phi_{v,t}$  is an automorphism. Moreover, since  $\phi_{v,t}$  does not preserve the trace function if  $t \neq 0$ , it is easy to see that  $\phi_{v,t}$  is not an inner automorphism unless  $v = t = 0$ , in which case it is the identity.

If  $\varphi$  is an automorphism of  $\mathrm{GL}_2(\mathbf{Z}/l^n)$ , then by [9, Cor. 2] the restriction of  $\varphi$  to  $\mathrm{SL}_2(\mathbf{Z}/l^n)$  coincides with that of the composite of some  $\phi_{v,t}$  with an inner automorphism. Applying the last part of Lemma 5 and noting that  $r_{\phi_{v,t,1,h}}$  is the identity map for any  $v$  and  $t$ , we have now determined the automorphisms of  $\mathrm{GL}_2(\mathbf{Z}/l^n)$ .

Finally, since  $l > 2$ , the trace zero elements of  $\mathrm{GL}_2(\mathbf{Z}/l^n)$  are precisely those whose squares are scalar matrices. Thus, they are taken to themselves under any automorphism, as desired. ■

#### 4. TWISTED MODULAR CURVES

In this section, we fix a positive integer  $N \geq 3$  and let  $S$  be a  $\mathbf{Z}[1/N]$ -scheme. Denote by  $\mathbf{Sch}_S$  and  $\mathbf{Sets}$  the category of  $S$ -schemes and sets, respectively. We define the (open) modular curve  $Y(N)$  over  $S$  as in [5, Cor. 4.7.2]. For any  $S$ -scheme  $T$ , we will denote  $Y(N) \times_S T$  by  $Y(N)$  when  $T$  is understood from context.

Given an elliptic curve  $E$  over a  $S$ -scheme  $T$ , denote by  $E[N]$  the  $N$ -torsion subgroup scheme of  $E$ . Since  $N$  is invertible over  $S$ , the finite locally free commutative group scheme  $E[N]$  is étale over  $T$  and after a finite étale surjective base change is isomorphic to the constant group scheme  $(\mathbf{Z}/N)^2$ . For any finite étale commutative group scheme  $G$  over  $S$  which is étale locally isomorphic to the constant group scheme  $(\mathbf{Z}/N)^2$ , we denote by  $\det G$  the finite étale  $S$ -group scheme which represents the étale sheaf  $\wedge_{\mathbf{Z}/N}^2(G)$ .

The following result is well-known to experts, but for the sake of completeness (and to assist the non-expert reader), we give a proof via reduction to standard results which are completely proven in [5].

**THEOREM 2.** *Let  $S$  and  $G$  be as above. For  $N \geq 3$ , the functor  $F_G: \mathbf{Sch}_S \rightarrow \mathbf{Sets}$  given by*

$$T \mapsto \left\{ \begin{array}{l} \text{isomorphism classes of pairs } (E, \alpha), \text{ with } E_{/T} \text{ an elliptic curve} \\ \text{and } \alpha: E[N] \simeq G \times_S T \text{ an isomorphism of } T\text{-group schemes} \end{array} \right\}$$

*is represented by an  $S$ -scheme  $Y(G)$  which becomes isomorphic to  $Y(N)$  over a finite étale cover of  $S$  (so  $Y(G) \rightarrow S$  is smooth and affine of pure relative dimension 1).*

Suppose we are given an isomorphism of  $S$ -group schemes  $i: \det G \simeq \mu_N$ . Then for  $N \geq 3$ , the functor  $F_G^i: \mathbf{Sch}_{/S} \rightarrow \mathbf{Sets}$  given by

$$T \mapsto \left\{ \begin{array}{l} \text{isomorphism classes of pairs } (E, \alpha), \text{ such that } E_{/T} \text{ is an elliptic} \\ \text{curve, } \alpha: E[N] \simeq G \times_S T \text{ is an isomorphism of } T\text{-group schemes,} \\ \text{and } \det \alpha: \det E[N] \simeq (\det G) \times_S T \simeq {}^i\mu_{N,T} \text{ is the Weil pairing} \end{array} \right\}$$

is represented by an open and closed subscheme  $Y(G, i)$  in  $Y(G)$ , and  $Y(G)$  is covered by the disjoint open subschemes  $Y(G, i^n)$  for  $n \in (\mathbf{Z}/N)^\times$ , where the isomorphism  $i^n$  is the composite of  $i$  and the  $n$ th power map on  $\mu_N$ . The scheme  $Y(G, i)$  has geometrically connected fibers over  $S$ .

*Proof.* We begin by showing that the functor  $F_G$  on  $\mathbf{Sch}_{/S}$  is an étale sheaf. Since  $F_G$  is trivially a Zariski sheaf (due to the rigidity of level  $N$  structures for  $N \geq 3$  [5, Cor. 2.7.2]), it remains to show that if  $T' \rightarrow T$  is a quasi-compact étale surjective map of  $S$ -schemes, then the diagram of sets

$$F_G(T) \rightarrow F_G(T') \rightrightarrows F_G(T' \times_T T') \tag{8}$$

is exact. Indeed, once such exactness is proven we can use étale descent theory to see that the representability of  $F_G$  by an affine smooth  $S$ -scheme with pure relative dimension 1 can be checked after we make a finite étale surjective base change  $S' \rightarrow S$  (the effectiveness of the descent data on affine  $S'$ -schemes with respect to  $S' \rightarrow S$  follows from [4, Cor. 7.6, Exp. VIII]). We can find such a base change so that  $G \times_S S' \simeq (\underline{\mathbf{Z}}/N)^2$ , so the representability over  $S'$  by the affine smooth  $S'$ -scheme  $Y(\overline{N})$  with pure relative dimension 1 follows from [5, Cor. 4.7.2].

By the rigidity of level  $N$  structures for  $N \geq 3$ ,  $F_G(T) \rightarrow F_G(T')$  is injective. Indeed, if  $(E_1, \alpha_1), (E_2, \alpha_2)$  over  $T$  become isomorphic over  $T'$ , via an isomorphism  $\varphi': E_1 \simeq E_2$  over  $T'$  that takes  $\alpha'_1$  to  $\alpha'_2$ , then both pullbacks of  $\varphi'$  to  $T'' = T' \times_T T'$  take  $\alpha''_1$  to  $\alpha''_2$ . By rigidity, we conclude that the two pullbacks of  $\varphi'$  to  $T' \times_T T'$  coincide, so by fpqc descent of morphisms we have  $\varphi' = \varphi \times_T T'$  for a unique map  $\varphi: E_1 \rightarrow E_2$  which is necessarily an isomorphism of elliptic curves taking  $\alpha_1$  to  $\alpha_2$ , as these properties all hold after the fpqc base change  $T' \rightarrow T$ . This establishes injectivity on the left of (8).

Now suppose that for some  $(E', \alpha')$  in  $F_G(T')$  there is an isomorphism  $\varphi: (E_1, \alpha_1) \simeq (E_2, \alpha_2)$  over  $T'' = T' \times_T T'$ , where  $(E_i, \alpha_i)$  is the base change by the  $i$ th projection  $T'' \rightarrow T'$ . We want to construct an  $(E, \alpha)$  in  $F_G(T)$  inducing  $(E', \alpha')$  in  $F_G(T')$ . By descent of schemes (using canonical projectiveness of elliptic curves to get effectiveness of descent data [4, Prop. 7.8, Exp. VIII]), it suffices to check that  $\varphi$  satisfies a ‘‘cocycle’’ condition. But this condition over  $T' \times_T T' \times_T T'$  is forced by the rigidity of level  $N$  structures for  $N \geq 3$ . This yields the desired exactness, so  $F_G$  is indeed an étale

sheaf on  $\mathbf{Sch}_{/S}$ . As we noted above, this implies the first part of the theorem, via reduction to the special case  $G = (\mathbf{Z}/N)^2$ .

To prove the second part of the theorem, denote by  $E^{\text{univ}} \rightarrow Y(G)$  the universal elliptic curve over  $Y(G)$ . The Weil pairing and  $\det \alpha$  give rise to a composite isomorphism

$$j: \mu_N \simeq \det E^{\text{univ}}[N] \simeq \det(G) \stackrel{i}{\simeq} \mu_N$$

over  $Y(G)$ , which is an automorphism  $\mu_N$  over  $Y(G)$ . Since  $\underline{\text{Aut}}(\mu_N) \simeq (\mathbf{Z}/N)^\times$  as étale sheaves on  $\mathbf{Sch}_{/S}$ ,  $j$  must be given Zariski locally on  $Y(G)$  by raising to the  $d$ th power for various  $d \in (\mathbf{Z}/N)^\times$ . It is obvious that  $F^{i^n}$  is represented by the open and closed subscheme  $Y(G, i^n)$  corresponding to  $d=n$ , and as  $n$  runs through the elements of  $(\mathbf{Z}/N)^\times$ , the  $Y(G, i^n)$ 's give a covering of  $Y(G)$  by disjoint open subschemes. Passing to geometric fibers, we may study the geometric connectedness of the fibers in the case  $S = \text{Spec } k$ , with  $k$  an algebraically closed field of characteristic not dividing  $N$  and  $G = (\mathbf{Z}/N)^2$ . In this case,  $\det G \simeq^i \mu_N$  corresponds to a choice of primitive  $N$ th root of unity  $\zeta_N \in \mu_N(k)$ . This choice makes  $k$  a  $\mathbf{Z}[1/N, \zeta_N]$ -algebra and  $Y(G, i)$  is exactly the  $k$ -fiber of the  $\mathbf{Z}[1/N, \zeta_N]$ -scheme  $Y(N)^{\text{can}}$  as defined in [5, 9.1.6]. However, it follows from [5, 10.9.2(2)] (which makes essential use of the complex analytic theory of modular curves and its compatibility with the algebraic theory) that  $Y(N)^{\text{can}}$  has geometrically connected fibers over  $\mathbf{Z}[1/N, \zeta_N]$ . ■

## 5. PROOF OF THEOREM 1

Let  $n > 1$  and choose a prime  $l = 3$  or  $5$ . Fix a number field  $K$  which is linearly disjoint from  $\mathbf{Q}(\zeta_{l^n})$ . Choose any  $r \in (\mathbf{Z}/l)^\times$  which is not a square. Let  $\mathcal{O}$  be the integer ring of  $K$ . By the Chebotarev density theorem and the linear disjointness of  $K$  and  $\mathbf{Q}(\zeta_l)$ , there exist infinitely many primes  $p \neq l$  in  $\mathbf{Z}$  such that  $p$  is totally split in  $K$  and  $p \equiv -r \pmod{l}$ . Fix a choice of such a  $p$ . In particular,  $X^2 + p$  does not have a root in the finite field  $\mathbf{F}_l$ . By Honda–Tate theory [8], there exists an elliptic curve  $\bar{E}_p$  over  $\mathbf{F}_p$  which is supersingular, which is to say that the characteristic polynomial of the arithmetic Frobenius action on the  $l$ -adic Tate module of  $\bar{E}_p$  is  $X^2 + p$ . Fix a choice of such a  $\bar{E}_p$  and choose a Weierstrass model for this over  $\mathbf{F}_p$ . Pick a prime  $\mathfrak{p}$  of  $K$  over  $p$  and choose a Weierstrass equation over  $\mathcal{O}_{\mathfrak{p}}$  whose reduction is the equation for  $\bar{E}_p$ . This defines an elliptic curve  $E_1$  over  $\mathcal{O}_{\mathfrak{p}}$  with reduction at  $\mathfrak{p}$  isomorphic to  $\bar{E}_p$ . Thus, the  $G_K$ -module action on  $E_1[l](\bar{K})$  must be irreducible, since  $X^2 + p$  has no roots in  $\mathbf{F}_l$ , and the same holds for any elliptic curve over  $K$  given by a Weierstrass equation which is  $\mathfrak{p}$ -adically close to that of  $E_1$ .

Choose any prime  $q$  of  $K$  not equal to  $p$  and not lying over  $l$ . From the theory of Tate curves [7, Ch. V, Thm. 5.3], we can find a Weierstrass equation over  $K$  which defines an elliptic curve  $E_2$  over  $K$  with split multiplication reduction at  $q$  and  $\text{ord}_q(j(E_2)) = -1$ . Moreover, any Weierstrass equation over  $K$  which is  $q$ -adically close to that of  $E_2$  will also have these properties. Now consider any elliptic curve  $E/K$  defined by a Weierstrass equation which is  $p$ -adically close to  $E_1$  and  $q$ -adically close to  $E_2$ . Clearly there are infinitely many  $j$ -invariant values  $j(E) \in K$  which arise in this way, and (by weak approximation) we can even find such  $E$  with good reduction at any desired finite set of places away from  $q$ , and split multiplicative reduction at  $q'$  with  $\text{ord}_{q'}(j(E)) = -1$  for any desired finite set of other places  $q'$  away from  $p$ . In particular, we can find an infinite set of such  $E$ 's so that the sets of ramified primes in the mod- $l^n$  representations of  $G_K$  are non-empty and mutually disjoint away from  $l$ .

We claim that the representation  $\rho_{E, l^n}: G_K \rightarrow \text{Aut}(E[l^n](\bar{K})) \simeq \text{GL}_2(\mathbf{Z}/l^n)$  is surjective for all such  $E$ . Since  $K$  is linearly disjoint from  $\mathbf{Q}(\zeta_{l^n})$ , it suffices to prove that  $\text{SL}_2(\mathbf{Z}/l^n)$  lies in the image of  $\rho_{E, l^n}$ . From the Tate parameterization of elliptic curves with split multiplicative reduction and the condition  $\text{ord}_q(j(E)) = -1$ , there is a basis  $\{e_1, e_2\}$  of  $E[l^n](\bar{K})$  over  $\mathbf{Z}/l^n$  with respect to which  $\sigma = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$  lies in the image of  $\rho_{E, l^n}$  on the inertia group at  $q$ . Since  $\rho_{E, l^n}(\text{mod } l)$  is irreducible, there exists  $g \in G_K$  such that  $e'_2 = ge_1 \notin (\mathbf{Z}/l^n)e_1$ . With respect to the basis  $\{e_1, e'_2\}$ , the automorphism  $\sigma$  becomes  $\begin{pmatrix} 1 & \alpha \\ & 1 \end{pmatrix}$ , whence  $g\sigma g^{-1} = \begin{pmatrix} 1 & \\ & \beta \end{pmatrix}$ , with  $\alpha, \beta \in \mathbf{Z}/l^n$ . Since  $\begin{pmatrix} 1 & \alpha \\ & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & \\ & \beta \end{pmatrix}$  are conjugate to  $\begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$  in  $\text{GL}_2(\mathbf{Z}/l^n)$ , these matrices have order  $l^n$ . Consequently, the image of  $\rho_{E, l^n}$  contains  $\begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & \\ & \beta \end{pmatrix}$ , which generate  $\text{SL}_2(\mathbf{Z}/l^n)$ . Thus the image of  $\rho_{E, l^n}$  contains  $\text{SL}_2(\mathbf{Z}/l^n)$ , so the representation  $\rho_{E, l^n}$  is surjective, as desired. Fix such an  $E$  as above and choose a basis of  $E[l^n](\bar{K})$  over  $\mathbf{Z}/l^n$ . Let  $\rho = \rho_{E, l^n}: G_K \rightarrow \text{GL}_2(\mathbf{Z}/l^n)$  be the corresponding representation.

Let  $\phi = \phi_{v, t}$  be an automorphism of  $\text{GL}_2(\mathbf{Z}/l^n)$  as furnished by Lemma 6 with  $v, t \not\equiv 0 \pmod{l^n}$ . Define  $\rho' = \phi \circ \rho$ , and let  $\bar{\rho}'$  be the induced mod- $l$  representation. Note that by the definition of  $\phi_{v, t}$ , the mod- $l$  representation  $\bar{\rho}$  obtained from  $\rho$  is literally equal to  $\bar{\rho}'$ . However,  $\rho$  and  $\rho'$  are *not* equivalent up to a twist. To see this, we note that if  $\rho$  and  $\rho'$  were equivalent up to a twist, then the corresponding projective representations would be conjugate. Since  $\rho$  is surjective and  $\rho' = \phi_{v, t} \circ \rho$ , it would follow that  $\phi_{v, t}$  induces an inner automorphism of  $\text{PGL}_2(\mathbf{Z}/l^n)$ , a contradiction (due to our choices of  $v$  and  $t$ ).

Viewing  $\rho'$  and  $\bar{\rho}'$  as finite étale group schemes over  $K$  with cyclotomic determinant, we denote by  $X(\rho')$  and  $X(\bar{\rho}')$  the canonical compactifications of the smooth affine curves as furnished by the first part of Theorem 2. There is an obvious natural  $K$ -morphism  $\pi: X(\rho') \rightarrow X(\bar{\rho}')$  which corresponds (away from the cuspidal part) to “reduction mod  $l$ ” in terms of

Yoneda's lemma. By the second part of Theorem 2, the connected components of  $X(\rho')$  and  $X(\bar{\rho}')$  are geometrically connected over  $K$ . We claim that the induced maps between connected components have degree  $l^{3(n-1)}$  (and in particular,  $\pi$  is finite flat). This can be checked after base change to  $\bar{K}$ , over which  $\pi$  becomes the canonical map  $X(l^n) \times_{\mathbf{Z}[1/l]} \bar{K} \rightarrow X(l) \times_{\mathbf{Z}[1/l]} \bar{K}$ , which is well-known to be a generically Galois covering between connected components, with Galois group  $\ker(\mathrm{PSL}_2(\mathbf{Z}/l^n) \rightarrow \mathrm{PSL}_2(\mathbf{Z}/l))$  having order  $l^{3(n-1)}$  (moreover, the branch locus is supported in the *cuspidal part*).

Since  $l=3$  or  $l=5$  and the genus of a proper smooth geometrically connected curve over a field can be computed after arbitrary change of the base field, the connected components of the proper smooth  $K$ -curve  $X(\bar{\rho}') = X(\bar{\rho})$  have genus 0. We claim that each of these connected components is  $K$ -isomorphic to  $\mathbf{P}_K^1$ . Let  $X$  be one of the connected components of  $X(\bar{\rho}') = X(\bar{\rho})$ , so  $X$  is a proper smooth *geometrically connected* curve over  $K$  with genus 0. In order to show that  $X \simeq \mathbf{P}_K^1$ , it suffices to show that  $X(K)$  is non-empty. There is a connected component  $X_1$  of  $X(\bar{\rho}') = X(\bar{\rho})$  which contains a  $K$ -rational point corresponding to the given elliptic curve  $E$  over  $K$  and the identity of  $G_K$ -modules  $E[l](\bar{K}) = \bar{\rho}$ . Since  $X_1(K)$  is non-empty,  $X_1 \simeq \mathbf{P}_K^1$ . It suffices below to just work with this component, but we want to briefly explain Mazur's elegant proof of the stronger result that all connected components of  $X(\bar{\rho}')$  are  $K$ -isomorphic to  $\mathbf{P}_K^1$ .

We see from the proof of Theorem 2 that the connected components of  $X(\bar{\rho})$  are indexed by elements  $v$  of  $(\mathbf{Z}/l)^\times$  (i.e., automorphisms of  $\mu_l$ ), and there is an obvious  $K$ -isomorphism of connected components  $X_v \simeq X_{vw^2}$  for any two  $v, w \in (\mathbf{Z}/l)^\times$ , by using Yoneda's Lemma and "multiplication by  $w$ " on the level of  $l$ -torsion group schemes. Thus, to show that all connected components of  $X(\bar{\rho}') = X(\bar{\rho})$  are  $K$ -isomorphic to  $\mathbf{P}_K^1$ , it suffices to show that  $X_v \simeq \mathbf{P}_K^1$  for a single non-square  $v \in (\mathbf{Z}/l)^\times$ . Since  $l=3$  or  $l=5$ , we may consider  $v=2$ . It is a classical observation that in order to show  $X(K)$  is non-empty, it suffices to construct a divisor  $D$  on  $X$  with odd degree. Indeed, adding a suitable multiple of the canonical divisor (which has degree  $-2$ ) to  $D$  gives a divisor  $D'$  on  $X$  with degree 1. By the Riemann–Roch Theorem for the *geometrically connected* proper smooth curve  $X$  over  $K$ , we have  $H^0(X, \mathcal{L}(D')) = 1 > 0$ , so there is an effective divisor on  $X$  with degree 1, which is to say that  $X(K)$  is non-empty. Thus, it suffices to construct a divisor with odd degree on  $X_2$ . Mazur observed that an étale-twisted version of the Hecke operator  $T_2$  gives a correspondence between  $X_1$  and  $X_2$  with degree 3 over both  $X_1$  and  $X_2$ . By using this correspondence and the existence of a  $K$ -rational point on  $X_1$ , we can construct an effective divisor on  $X_2$  with odd degree (1 or 3). This completes the sketch of Mazur's proof that every connected component of  $X(\bar{\rho}')$  is  $K$ -isomorphic to  $\mathbf{P}_K^1$ .

Fix a connected component  $C \simeq \mathbf{P}_K^1$  of  $X(\bar{\rho}')$  and a connected component  $C'$  in  $X(\rho')$  over  $C$ , so  $\pi_{C'}: C' \rightarrow C$  is a finite map with degree  $l^{3(n-1)}$ . By Theorem 2, the proper smooth curve  $C'$  over  $K$  is geometrically connected. Therefore, by Theorem 1, there exist infinitely many non-cuspidal  $a \in C(K)$  such that  $\pi_{C'}^{-1}(a) = \text{Spec}(L_a)$ , where  $L_a$  is a finite extension of  $K$  which is linearly disjoint from the splitting field of  $\rho$  (which coincides with the splitting field of  $\rho'$ ). Obviously  $[L_a:K]$  is equal to the degree of  $\pi_{C'}$ , which is  $l^{3(n-1)}$ . From the linear disjointness, it follows that the representations  $\rho|_{G_{L_a}}$  and  $\rho'|_{G_{L_a}}$  are surjective and come from the mod- $l^n$  representations of elliptic curves over  $L_a$  with  $j$ -invariants in  $K$  (since  $a \in C(K)$ ). Of course, we can choose these  $j$ -invariants to avoid any desired finite set of elements of  $K$ . By the choice of  $\phi$ ,  $\rho|_{G_{L_a}}$  and  $\rho'|_{G_{L_a}}$  satisfy the condition (1) in the Introduction and are not equivalent up to twists.

It remains to analyze ramification in  $L_a/K$ . Recall that in our construction of elliptic curves above via Tate models, we saw that we can choose the mod- $l^n$  representation  $\rho$  coming from our elliptic curve  $E$  over  $K$  to be ramified at any desired finite set of primes of  $K$  away from  $l$  and to be unramified at any desired finite set of other primes of  $K$  away from  $l$ . In order to complete the proof of the theorem, we need to check that the ramification in  $L_a$  outside of  $l$  can be chosen to avoid any desired finite set of primes of  $K$  with norm  $> (l^2 - 3)/2$ .

Choose a prime  $\mathfrak{p}$  of  $K$  not over  $l$  at which  $E$  has good reduction. Thus,  $\rho$  and  $\rho'$  are unramified at  $\mathfrak{p}$ . By étale descent, we can identify  $\rho'$  with the generic fiber of a finite étale group scheme  $\mathcal{G}$  over  $\mathcal{O}_{\mathfrak{p}}$  which is étale-locally isomorphic to the constant group scheme  $(\mathbf{Z}/l^n)^2$ . The  $l$ -torsion subgroupscheme  $\mathcal{G}[l] \simeq \mathcal{G}/l^{n-1}$  is an analogous  $\mathcal{O}_{\mathfrak{p}}$ -model for  $\bar{\rho}'$ . Since  $\mathcal{O}_{\mathfrak{p}}$  is a  $\mathbf{Z}[1/l]$ -scheme, we can use Theorem 2 and the compactification theory of modular curves [5, 8.6.7, 10.9.5] to realize the map  $\pi: X(\rho') \rightarrow X(\bar{\rho}')$  as the  $K$ -fiber of a finite flat map  $\pi_{\mathfrak{p}}: X(\mathcal{G}) \rightarrow X(\mathcal{G}[l])$  between proper smooth  $\mathcal{O}_{\mathfrak{p}}$ -schemes with geometric fibers of pure dimension 1. This map  $\pi_{\mathfrak{p}}$  is just an étale twist of the finite flat map  $X(l^n) \times_{\mathbf{Z}[1/l]} \mathcal{O}_{\mathfrak{p}} \rightarrow X(l) \times_{\mathbf{Z}[1/l]} \mathcal{O}_{\mathfrak{p}}$ .

Since the natural map  $X(l^n) \rightarrow X(l)$  over  $\mathbf{Z}[1/l]$  is Galois away from the cusps, the branch locus of  $\pi_{\mathfrak{p}}$  is supported in the cuspidal subscheme of  $X(\mathcal{G}[l])$ , which is étale over  $\mathcal{O}_{\mathfrak{p}}$  with degree  $(l^2 - 1)/2$  (as the same holds for the cuspidal subscheme of  $X(l)$  over  $\mathbf{Z}[1/l]$ ). Consider  $a \in C(K) \subseteq X(\bar{\rho}')(K) = X(\mathcal{G}[l])(K)$  as above. The scheme-theoretic closure of  $a \in X(\mathcal{G}[l])(K)$  in  $X(\mathcal{G}[l])$  is a point  $\bar{a} \in X(\mathcal{G}[l])(\mathcal{O}_{\mathfrak{p}})$ , by the valuative criterion for properness. If we can choose  $a$  so that the closed point of  $\bar{a}$  is not a cusp, then  $\pi_{C'}^{-1}(a) = \text{Spec}(L_a)$  is a component of the generic fiber of the finite étale scheme  $\pi_{\mathfrak{p}}^{-1}(\bar{a})$  over  $\bar{a} = \text{Spec}(\mathcal{O}_{\mathfrak{p}})$ . Thus, the prime  $\mathfrak{p}$  would not ramify in  $L_a$ . In order to check that  $a$  can be chosen in the manner desired, consider the connected component  $\bar{C}$  of  $X(\mathcal{G}[l])$  which



has generic fiber  $C$  (so  $\bar{a} \in \bar{C}(\mathcal{O}_{\mathfrak{p}})$ ). Since  $\bar{C} \rightarrow \text{Spec}(\mathcal{O}_{\mathfrak{p}})$  is proper and smooth with geometric fibers of pure dimension 1 and generic fiber  $\mathbf{P}_K^1$ , it must be the case that  $\bar{C} \simeq \mathbf{P}_{\mathcal{O}_{\mathfrak{p}}}^1$ , thanks to the following well-known lemma. We give a proof due to lack of an adequate reference.

**LEMMA 7.** *Let  $R$  be a discrete valuation ring with fraction field  $K$ ,  $X$  a proper smooth  $R$ -scheme with pure relative dimension 1 and generic fiber  $X \times_R K \simeq \mathbf{P}_K^1$ . Then  $X \simeq \mathbf{P}_R^1$  over  $R$ .*

*Proof.* Since the generic fiber of  $X$  is geometrically connected, the closed fiber is also geometrically connected [3, IV<sub>3</sub>, 12.2.4(vi)], necessarily with genus 0. By the valuative criterion for properness, we have  $X(R) = X(K)$ . This set is non-empty, so choose a section  $\text{Spec}(R) \rightarrow X$  over  $R$ . This defines a relative effective Cartier divisor  $D$  on  $X$  over  $R$  with degree 1. By Grothendieck's theory of cohomology and base change, as well as the Riemann–Roch theorem for genus 0 curves over fields,  $\mathcal{L}(D)$  is generated by its global sections  $H^0(X, \mathcal{L}(D))$  and this  $R$ -module is locally free of rank 2 over  $R$ , commuting with arbitrary base change over  $R$ . Since  $R$  is local,  $H^0(X, \mathcal{L}(D))$  is free of rank 2. Choosing a basis gives a map  $X \rightarrow \mathbf{P}_R^1$  which commutes with arbitrary base change over  $R$ . We claim this map is an isomorphism. Since both sides are smooth over  $R$ , by [3, IV<sub>4</sub>, 17.9.5] it suffices to show that the induced map on fibers over  $\text{Spec}(R)$  is an isomorphism. But over a field  $k$ , it is classical that for a proper, smooth, geometrically connected curve  $C$  over  $k$  with genus 0, and a rational function  $f \in k(C)$  with a simple pole at a  $k$ -rational point and no other poles, the map  $f: C \rightarrow \mathbf{P}_k^1$  is an isomorphism. ■

Thus, as long as the number of rational points  $|\mathcal{O}/\mathfrak{p}| + 1$  in the closed fiber of  $\bar{C} \simeq \mathbf{P}_{\mathcal{O}_{\mathfrak{p}}}^1$  is larger than the degree  $(l^2 - 1)/2$  ( $=4$  or  $12$ ) of the cuspidal subscheme on  $X(\mathcal{G}[l])$ , then a  $\mathfrak{p}$ -adic congruence condition on  $a \in C(K) = \mathbf{P}_K^1$  ensures that the closed point of  $\bar{a}$  is non-cuspidal. This implies that  $\bar{a}$  is disjoint from the branch locus of  $\pi_{\mathfrak{p}}$ , so  $L_a$  is unramified over  $\mathfrak{p}$ . Thus, we can indeed force  $\mathfrak{p}$  to be unramified in  $L_a$  if the norm of  $\mathfrak{p}$  exceeds  $(l^2 - 3)/2$ . The same argument allows us to handle any finite number of such  $\mathfrak{p}$ 's simultaneously.

## ACKNOWLEDGMENTS

S.W. thanks Karl Rubin for a useful discussion on twisted modular curves. Both authors thank the referee for helpful comments.

## REFERENCES

1. P. M. Cohn, On the structure of the  $GL_2$  of a ring, *Publ. Math. IHES* **30** (1966), 5–54.
2. M. H. Dull, Automorphisms of the two-dimensional linear groups over integral domains, *Amer. J. Math.* **96** (1974), 1–40.
3. A. Grothendieck, Éléments de géométrie algébrique, *Publ. Math. IHES* **24**, **28**, **32** (1966).
4. A. Grothendieck, “Revêtements étales et groupe fondamental,” *Lecture Notes in Mathematics*, Vol. 224, Springer-Verlag, New York, 1971.
5. N. Katz and B. Mazur, “Arithmetic Moduli of Elliptic Curves,” Princeton Univ. Press, Princeton, NJ, 1985.
6. S. Lang, “Fundamentals of Diophantine Geometry,” Springer-Verlag, Berlin/New York, 1983.
7. J. Silverman, “Advanced Topics in the Arithmetic of Elliptic Curves,” Springer-Verlag, New York/Berlin, 1994.
8. J. Tate, Classes d’isogénie des variétés abéliennes sur un corps fini (d’après T. Honda), *Sém. Bourbaki* **352** (1968).
9. S. Wong, Twists of Galois representations and projective automorphisms, *J. Number Theory*, to appear.