

## Math 52H: Practice problems for the midterm

1 Consider the Euclidean space  $V = \mathbb{R}^{2n}$  with coordinates  $(x_1, y_1, \dots, x_n, y_n)$  and the standard dot-product. The space  $\Lambda(V^*)$  of all exterior  $k$ -forms for all  $k = 0, \dots, 2n$  is also an Euclidean space with the scalar product of a  $k$ -form  $\alpha$  and an  $l$ -form  $\beta$  defined by the formula

$$\langle\langle \alpha, \beta \rangle\rangle = \begin{cases} \star^{-1}(\alpha \wedge \star\beta), & \text{if } k = l, \\ 0, & \text{if } k \neq l. \end{cases}$$

Consider a linear operator  $\Omega : \Lambda(V^*) \rightarrow \Lambda(V^*)$  defined by the formula  $\Omega(\alpha) = \alpha \wedge \omega$ , where  $\omega = \sum_1^n x_i \wedge y_1$ . Find the adjoint linear operator  $\Omega^*$ , i.e. the operator  $\Omega^* : \Lambda(V^*) \rightarrow \Lambda(V^*)$  such that

$$\langle\langle \Omega(\alpha), \beta \rangle\rangle = \langle\langle \alpha, \Omega^*(\beta) \rangle\rangle$$

for any forms  $\alpha, \beta \in \Lambda(V^*)$ .

2. The cylindrical coordinates

$$r \in [0, \infty), \varphi \in [0, 2\pi), z \in \mathbb{R},$$

are introduced in  $\mathbb{R}^3$  by the formulas

$$x = r \cos \varphi, y = r \sin \varphi, z,$$

where  $(x, y, z)$  are Cartesian coordinates. Consider a differential 1-form

$$\alpha = \cos r dz + \frac{r \sin r}{\pi} d\varphi.$$

a) Describe the plane field  $\xi$  defined by the Pfaffian equation  $\alpha = 0$ .

b) Suppose that a curve  $\Gamma \subset \mathbb{R}^3$  be given by the parametric equations

$$r = \frac{\pi}{4}, z = h(t), \varphi = 2t, t \in [0, \pi].$$

Find the function  $h$  such that  $\alpha|_{\Gamma} = 0$  and  $h(0) = 1$ .

c) Find all values of  $R$  for which the horizontal circles  $\{z = \text{const}, r = R\}$  are tangent to the plane field  $\xi$ .

3. Consider a smooth function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ . Let  $S_f$  be a surface in  $\mathbb{R}^4$  given by equations

$$x_3 = \frac{\partial f}{\partial x_1}(x_1, x_2), \quad x_4 = \frac{\partial f}{\partial x_2}(x_1, x_2) \quad (1)$$

Suppose that this system of equations can be solved with respect to the coordinates  $x_2$  and  $x_4$ , i.e. there exist smooth functions  $x_2 = g(x_1, x_3)$  and  $x_4 = h(x_1, x_3)$  such that

$$\begin{aligned} x_3 &\equiv \frac{\partial f}{\partial x_1}(x_1, g(x_1, x_3)), \\ h(x_1, x_3) &\equiv \frac{\partial f}{\partial x_2}(x_1, g(x_1, x_3)). \end{aligned} \quad (2)$$

Prove that the Jacobian of the map  $(h, g) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is equal to  $-1$ , i.e. that

$$\begin{vmatrix} \frac{\partial g}{\partial x_1} & \frac{\partial g}{\partial x_3} \\ \frac{\partial h}{\partial x_1} & \frac{\partial h}{\partial x_3} \end{vmatrix} = -1.$$

Hint: Examine the restriction of the form  $\omega = dx_1 \wedge dx_3 + dx_2 \wedge dx_4$  to the surface  $S_f$ , and then consider the pull-back of the form  $\omega$  by a map  $\mathbb{R}^2 \rightarrow S_f \subset \mathbb{R}^4$  given by the formulas

$$(x_1, x_3) \mapsto (x_1, g(x_1, x_3), x_3, h(x_1, x_3)).$$

4. Consider a differential 1-form  $\alpha = dx_3 + x_2 dx_1$  on  $\mathbb{R}^3$ . Let  $f = (f_1, f_2, f_3) : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a map such that  $f^* \alpha = h \alpha$  for some positive function  $h : \mathbb{R}^3 \rightarrow \mathbb{R}$ . Find a function  $g : \mathbb{R}^3 \rightarrow \mathbb{R}$  such that the map  $F : \mathbb{R}^4 \rightarrow \mathbb{R}^4$  given by the formula

$$F(x_1, x_2, x_3, x_4) = (f_1(x_1, x_2, x_3), f_2(x_1, x_2, x_3), f_3(x_1, x_2, x_3), x_4 g(x_1, x_2, x_3))$$

satisfies the equation  $F^*(x_4\alpha) = x_4\alpha$ .

5. Consider a smooth differential  $k$ -form

$$\alpha = \sum_{1 \leq i_1 < \dots < i_k \leq n} f_{i_1 \dots i_k}(x) dx_{i_1} \wedge \dots \wedge dx_{i_k}$$

in  $\mathbb{R}^n$  such that  $f_{i_1 \dots i_k}(0) = 0$  (i.e. all coefficients of the form  $\alpha$  are equal to 0 at the origin).

Let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  denote the dilatation  $x \mapsto 2x$ . Suppose that  $F^*\alpha = \alpha$ . Prove that  $\alpha \equiv 0$ .

6. Given a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , consider a map  $F : \mathbb{R}^n \rightarrow \mathbb{R}^{2n+1}$  defined by the formula

$$F(x_1, \dots, x_n) = \left( x_1, \dots, x_n, \frac{\partial f}{\partial x_1}(x_1, \dots, x_n), \dots, \frac{\partial f}{\partial x_n}(x_1, \dots, x_n), f(x_1, \dots, x_n) \right).$$

Compute  $F^*(\alpha)$ , where

$$\alpha = dx_{2n+1} - \sum_{i=1}^n x_{i+n} dx_i.$$

The actual midterm will consist of four problems.