

Math 52H: Homework N2

Due to Friday, January 25

1. Do Exercise 4.5 from the online text:

- (a) For any special orthogonal operator $\mathcal{A} : V \rightarrow V$ the operators $\mathcal{A}^* : \Lambda^k(V^*) \rightarrow \Lambda^k(V^*)$ and \star commute, i.e.

$$\mathcal{A}^* \circ \star = \star \circ \mathcal{A}^*.$$

- (b) Let A be an orthogonal matrix of order n with $\det A = 1$. Prove that the absolute value of each k -minor M of A is equal to the absolute value of its complementary minor of order $(n - k)$. Here k is any integer between 1 and n . (*Hint:* Apply (a) to the form $x_{i_1} \wedge \cdots \wedge x_{i_k}$).

2. Denote coordinates in \mathbb{R}^{2n} by $x_1, y_1, \dots, x_n, y_n$. Let ω denote the 2-form $\sum_1^n x_i \wedge y_i$.

A linear operator $\mathcal{A} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ is called *symplectic* if $\mathcal{A}^*\omega = \omega$.

Denote

$$J = \begin{pmatrix} 0 & -1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & -1 & \dots & 0 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ & & & & \dots & & \\ 0 & 0 & 0 & 0 & \dots & 0 & -1 \\ 0 & 0 & 0 & 0 & \dots & 1 & 0 \end{pmatrix}.$$

Note that $J^2 = -I$. Let $\mathcal{J} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ be the operator with the matrix J .

An operator $\mathcal{U} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ is called *unitary* if it is orthogonal and commutes with \mathcal{J} , i.e. $\mathcal{U} \circ \mathcal{J} = \mathcal{J} \circ \mathcal{U}$. Respectively, a matrix U is called unitary, if it is orthogonal and $UJ = JU$.

(b) Prove that an operator \mathcal{A} is symplectic if and only if its matrix A (in the standard basis of \mathbb{R}^{2n}) satisfies the equation $A^T J A = J$.

(c) Prove that an orthogonal operator $\mathcal{U} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ is unitary if and only if it is symplectic.

(d) An n -dimensional vector subspace $L \subset \mathbb{R}^{2n}$ is called *Lagrangian* if $\omega|_L = 0$. Prove that L is Lagrangian if and only if $\mathcal{J}(L) = L^\perp$.

3. A 2-form β on \mathbb{R}^4 is called self-dual if $\star\beta = \beta$. What is the dimension of the space of self-dual 2-forms on \mathbb{R}^4 . Find a basis of this space.

4. Consider $V = \mathbb{R}^3$ with the dot-product. Show that the formula

$$X \times Y = \mathcal{D}^{-1}(\star(\mathcal{D}(X) \wedge \mathcal{D}(Y)))$$

defines the cross-product on \mathbb{R}^3 .

Recall that the cross-product $X \times Y$ is defined as the vector Z orthogonal to $\text{Span}(X, Y)$ which has length equal to the area of the parallelogram $P(X, Y)$ and (assuming that X, Y are linearly independent) directed in such a way that the basis (X, Y, Z) defines the standard orientation of \mathbb{R}^3 . We also recall that the isomorphism $\mathcal{D} : V \rightarrow V^*$ is defined by the formula $\mathcal{D}(v)(X) = \langle v, X \rangle$, $v, X \in V$.

5.* (*Cartan's lemma*). Suppose that linear functions $l_1, \dots, l_k \in V^*$ are linearly independent and $\lambda_1, \dots, \lambda_k \in V^*$ are any other linear functions. Prove that

$$l_1 \wedge \lambda_1 + \dots + l_k \wedge \lambda_k = 0$$

if and only if

$$\lambda_1, \dots, \lambda_k \in \text{Span}(l_1, \dots, l_k)$$

and the matrix $A = (a_{ij})$ defined by $\lambda_i = \sum_{j=1}^k a_{ij} l_j$ is symmetric.

All problems and their subproblems are 10 points each. The problem 5 is an extra-credit.