MODERN ALGEBRA (MATH 210) PROBLEM SET 3

1. (a) Must any finite integral domain be a field? (b) *(The field $\mathbb{F}_4$.)* Ex. 3.14 (p. 125). Ex. 3.44 (ii) (p. 149).

2. Prove that the quotient ring $\mathbb{Z}[i]/I$ is finite for any nonzero ideal I of $\mathbb{Z}[i]$. (*Hint:* Use the fact that $I = (\alpha)$ for some nonzero α and then use the division algorithm to see that every coset of I is represented by an element of norm less than $N(\alpha)$.)

3.

(a) Prove that if an integer is the sum of two rational squares, then it is the sum of two integer squares (for example, $13 = (1/5)^2 + (18/5)^2 = 2^2 + 3^2$).
(b) Determine all the representations of the integer $2130797 = 17^2 \cdot 73 \cdot 101$ as the sum of two squares.
(c) Find a generator for the ideal $(47 - 13i, 53 + 56i)$ in $\mathbb{Z}[i]$.

4. Let I and J be ideals of a commutative ring R.

(a) Prove that $I + J$ is the smallest ideal of R containing both I and J.
(b) Prove that IJ is an ideal contained in $I \cap J$.
(c) Given an example where $IJ \neq I \cap J$.
(d) Prove that if $I + J = R$ then $IJ = I \cap J$.

5. A commutative ring R is called a *local ring* if it has a unique maximal ideal. Prove that if R is a local ring with maximal ideal $\mathfrak{m}$ then every element of $R - \mathfrak{m}$ is a unit. Prove conversely that if R is a commutative ring with 1 in which the set of nonunits forms an ideal $\mathfrak{m}$, then R is a local ring with unique maximal ideal $\mathfrak{m}$.

6. Let ω be the cube-root of unity $\frac{-1+\sqrt{-3}}{2}$. Show that the ring $R = \mathbb{Z}[\omega]$ (sometimes called the *Eisenstein integers*) has unique factorization. (*Hint:* see the proof for the Gaussian integers in Sec. 3.6.) Factor 2, 3, 5, and 7 into primes in R. (Which one of them has a repeated prime factor? This prime factor is key to an elementary proof of Fermat's Last Theorem for $n = 3$. If you'd like to see it, just ask me.)

7. Let K be a field. A *discrete valuation* on K is a function $v : K^\times \to \mathbb{Z}$ satisfying

(i) $v(ab) = v(a) + v(b)$ (i.e., v is a homomorphism from the multiplicative group of nonzero elements of K to $\mathbb{Z}$),
(ii) v is surjective, and
(iii) $v(x + y) \geq \min\{v(x), v(y)\}$ for all $x, y \in K^\times$ with $x + y \neq 0$.

Date: Tuesday, October 22, 2002.

The set $R = \{x \times K^\times : v(x) \geq 0\} \cup \{0\}$ is called the *valuation ring* of v.

(a) Prove that R is a subring of K which contains the identity. (In general, a ring R is called a *discrete valuation ring* if there is some field K and some discrete valuation v on K such that R is the valuation ring of v. In fact $K = \mathrm{Frac}(R)$.)
(b) Prove that for each nonzero element $x \in K$ either x or x^{-1} is in R.
(c) Prove that an element x is a unit of R if and only if $v(x) = 0$.
(d) *The p-adic valuation v_p.* Let p be a prime. One example of a discrete valuation is $v_p : \mathbb{Q}^\times \to \mathbb{Z}$ given by $v_p(a/b) = \alpha$ where $a/b = p^\alpha c/d$, with $(p, cd) = 1$. Describe the element of the corresponding valuation ring. Describe the units of the valuation ring.

Remark: There is a theorem that the only valuations on $\mathbb{Z}$ are these and the usual one. You can do analysis on $\mathbb{Q}$ using these valuations; they and their generalizations are central in number theory.

8. Let F be a field. Define the ring $F((x))$ of *formal Laurent series* with coefficients in F by

$$F((x)) = \left\{ \sum_{n \geq N}^{\infty} a_n x^n : a_n \in F, N \in \mathbb{Z} \right\}.$$

(Every element of $F((x))$ is a power series in x plus a polynomial in $1/x$, i.e. each element of $F((x))$ has only a finite number of terms with negative powers of x.)

(a) Prove that $F((x))$ is a field.
(b) Define a discrete valuation on $F((x))$ whose discrete valuation ring is $F[[x]]$, the ring of formal power series with coefficients in F.

Hence $F((x)) = \mathrm{Frac}(F[[x]])$. (This can be shown directly as well.)

9. Let D be a squarefree integer, and let $\mathcal{O} = \mathbb{Z}[\omega]$ be the ring of integers in the quadratic field $\mathbb{Q}(\sqrt{D})$, where $\omega = (1 + \sqrt{D})/2$ if $D \equiv 1 \pmod 4$, and $\omega = \sqrt{D}$ otherwise. For any positive integer f prove that the set $\mathcal{O}_f = \mathbb{Z}[f\omega] = \{a + bf\omega : a, b \in \mathbb{Z}\}$ is a subring of $\mathcal{O}$. Prove that $[\mathcal{O} : \mathcal{O}_f] = f$ (where the index is as additive abelian groups). Prove conversely that a subring of $\mathcal{O}$ containing the identity and having finite index f in $\mathcal{O}$ is equal to $\mathcal{O}_f$. (The ring $\mathcal{O}_f$ is called the *order of conductor f* in the field $\mathbb{Q}(\sqrt{D})$. The ring of integers $\mathcal{O}$ is called the *maximal order* in $\mathbb{Q}(\sqrt{D})$.)

10. *An example of a non-commutative ring.* Suppose G is a finite group. The *group ring* of G with coefficients in a commutative ring R, denoted RG, is defined on p. 521 (where it is called a *group algebra*). *Warning:* this is a non-commutative ring if G is not abelian!

1. Let $\mathcal{K} = \{k_1, \ldots, k_m\}$ be a conjugacy class in G. Prove that the element $K := k_1 + \cdots + k_m$ is in the center of RG.
2. Let $\mathcal{K}_1, \ldots, \mathcal{K}_r$ be the conjugacy classes of G, and for each $\mathcal{K}_i$ let K_i be the element of RG that is the sum of the members of $\mathcal{K}_i$. Prove that an element $\alpha \in RG$ is in the center of RG if and only if $\alpha = a_1 K_1 + a_2 K_2 + \cdots + a_r K_r$ for some $a_1, \ldots, a_r \in R$.

11. Let R be the ring of all continuous functions from $\mathbb{R}$ to $\mathbb{R}$, and for each $c \in \mathbb{R}$ let M_c be the ideal $\{f \in R : f(c) = 0\}$.

(a) Show that M_c is a maximal ideal.
(b) Let I be the collection of functions $f(x)$ in R with *compact support* (i.e. $f(x) = 0$ for $|x|$ sufficiently large) Prove that I is an ideal of R that is not a prime ideal.
(c) Let M be a maximal ideal of R containing I. Prove that $M \neq M_c$ for any $c \in \mathbb{R}$.

12. A universal side divisor u is a non-zero, non-unit element of R such that for every $x \in R$, either $u|x$ or $u|(x + z)$ for some unit z.

A Euclidean domain has universal side divisors: fix the norm N, and let u be an element of $R - \{0, \text{units}\}$ of minimal norm. Then for any $x \in R$, write $x = qu + r$ where $r = 0$ or $N(r) < N(u)$ (i.e. r is a unit). Thus u divides x in the first case, and $x - r$ in the second. (This argument is given in Prop. 3.63 on p. 154.)

Suppose $R = \mathbb{Z}[\alpha]$ where $\alpha = (1 + \sqrt{-19})/2$. Prove that R is not a Euclidean domain by showing that it has no universal side divisors as follows.

(a) Find the units of R.
(b) Take the usual norm N on R ($N(z) = z\overline{z}$). Show that the only elements of norm less than 5 are $\{0, \pm 1, \pm 2\}$.
(c) Taking $x = 2$ in the definition of a universal side divisor, show that any universal side divisor must be a nonunit divisor of 2 of 3.
(d) Find all nonunit divisors of 2 and 3.
(e) Now taking $x = \alpha$ in the definition of universal side divisor, find a contradiction.
(f) Repeat the argument for $\alpha = (-1 + \sqrt{-163})/2$.

It is not hard to show that $\mathbb{Q}(\alpha)$ is a Principal Ideal Domain, so this shows that not every Principal Ideal Domain is a Euclidean Domain.

The set is due Tuesday, October 29 at 3:30 pm in Pierre Albin's mailbox (opposite the elevator on the first floor of Building 380).