

# COMPLEX ALGEBRAIC SURFACES CLASS 16

RAVI VAKIL

## CONTENTS

1. Castelnuovo's Theorem	1
1.1. Motivation: Minimal rational surfaces	1
1.2. Motivation Luroth's theorem (in characteristic 0)	2
2. Proof of Castelnuovo's criterion (part 1)	2

On board beforehand:

- Useful trick.  $|D| \neq \emptyset$  (i.e.  $h^0(D) > 0$ ),  $C$  irreducible,  $C^2 \geq 0$  implies  $DC \geq 0$ .
- Genus formula.  $2g(C) - 2 = C \cdot (K_S + C)$ .
- Riemann-Roch:  $\chi(D) = \chi(\mathcal{O}) + \frac{1}{2}D \cdot (D - K)$ .
- Riemann-Roch: In the case when  $h^1(\mathcal{O}) = 0$  and  $h^0(K - D) = 0$ , we have  $h^0(D) \geq 1 + \frac{1}{2}D \cdot (D - K)$  (with equality iff  $h^1(D) = h^2(\mathcal{O}) = 0$ ).

## 1. CASTELNUOVO'S THEOREM

We saw how tricky it was to show that a surface is rational.

**Theorem: Castelnuovo's Rationality Criterion.** Let  $S$  be a surface with  $q = P_2 = 0$ . Then  $S$  is rational.

*Reminder.*  $q = h^1(S, \mathcal{O}_S) = h^0(S, \Omega_S) = h^2(S, \Omega_S) = h^1(S, K_S)$  (draw Hodge diamond). This is called the *irregularity* of a surface.

$$P_2 = h^0(S, K_S^{\otimes 2}).$$

It was once believed that this could be weakened to  $q = P_1 = 0$ , which is somehow more attractive (as  $P_1$  is an entry in the Hodge diamond), but this false, and we may see examples before the end of the course (Enriques surfaces, Godeaux surfaces).

**1.1. Motivation: Minimal rational surfaces.** We know lots of rational surfaces now:  $\mathbb{P}^2$ ,  $\mathbb{F}_n$ , and blow-ups of these. At this point, we may suspect that we've found them all. How can we show this? We'll use Castelnuovo's criterion.

---

*Date:* Friday, November 22.

**1.2. Motivation Luroth's theorem (in characteristic 0).** A variety  $V$  of dimension  $n$  is *unirational* if there is a dominant map (i.e. one with dense image)  $\mathbb{P}^n \dashrightarrow V$ .

**Lüroth's Theorem.** Every unirational curve is rational.

*Proof.* This is true in arbitrary characteristic, but here's a proof that works only in characteristic 0. Suppose  $\mathbb{P}^1 \dashrightarrow C$ , where  $C$  is a curve, possibly singular and not proper. Then we also get a rational map  $\mathbb{P}^1 \dashrightarrow C'$ , where  $C'$  is a smooth compactification of a smoothing of  $C$ . By our lemma from long ago, any rational map from a smooth curve to anything projective extends to a morphism, so we have  $\mathbb{P}^1 \rightarrow C'$ . Dominant implies surjective. So we can apply the Riemann-Hurwitz formula, to see that

$$2 - 2g(\mathbb{P}^1) = d(2 - 2g(C')) - \text{ramification contribution}.$$

The left side is 2, but if  $g(C') > 0$  the right side can't be positive. □

**Theorem.** In characteristic 0, every unirational surface is rational.

In positive characteristic, the theorem is *false*! Ask Ted Hwa for an example.

Question: where does the following argument break down in positive characteristic?

*Proof.* Suppose  $S$  is a unirational surface. If there was any doubt, let's say that it is smooth and compact. (Otherwise, there is a way of producing a smooth and compact birational model.) So we have  $\mathbb{P}^2 \dashrightarrow S$ . By the elimination of indeterminacy, we can blow up  $\mathbb{P}^2$  and get a morphism  $\text{Bl } \mathbb{P}^2 \rightarrow S$ . This morphism is dominant and hence surjective. Interpret  $q(S)$  as  $H^0(S, \Omega_S)$ , and recall  $P_2(S) = H^0(S, \mathcal{K}_S^{\otimes 2})$ . If  $q > 0$  or  $P_2 > 0$ , then pullback the nonzero form (i.e. section of either  $\Omega_S$  or  $\mathcal{K}_S^{\otimes 2}$ ) to get a non-zero section of the corresponding bundle on  $\text{Bl}(\mathbb{P}^2)$ . This would give  $q(\text{Bl}(\mathbb{P}^2)) > 0$  or  $P_2(\text{Bl}(\mathbb{P}^2)) > 0$ .

Hence  $q(S) = P_2(S) = 0$ . Then by Castelnuovo,  $S$  is rational. □

**Remark.** Even in characteristic 0, there are 3-folds that are unirational but not rational, and they are not even that exotic! It is not hard to show that smooth cubic threefolds in  $\mathbb{P}^4$  are all unirational; Clemens and Griffiths showed that *none* of them are rational! Iskovskih and Manin did the same for quartic threefolds as well.

## 2. PROOF OF CASTELNUOVO'S CRITERION (PART 1)

We'll make a couple of reduction steps.

**Castelnuovo'.** Let  $S$  be a minimal surface with  $q = P_2 = 0$ . Then there exists a smooth rational curve  $C$  on  $S$  such that  $C^2 \geq 0$ . *Keep on board.*

*Proof that Castelnuovo' implies Castelnuovo's criterion.*

$\mathcal{O}_S(C)$  clearly has a section, one whose zero set is  $C$ . We'll see that in fact  $h^0(S, \mathcal{O}_S(C)) \geq 2$ , so "the curve moves". Consider  $0 \rightarrow \mathcal{O}_S \rightarrow \mathcal{O}_S(C) \rightarrow \mathcal{O}_C(C) \rightarrow 0$ . Now  $q = h^1(S, \mathcal{O}_S) =$

0, so when we take global sections, the sequence remains exact, so

$$\begin{aligned}
h^0(S, \mathcal{O}_S(C)) &= h^0(S, \mathcal{O}_S) + h^0(C, \mathcal{O}_C(C)) \\
&= 1 + C^2 - g(C) + 1 + h^1(C, \mathcal{O}_C(C)) \\
&= 2 + C^2 \quad (\text{as } C \cong \mathbb{P}^1, \text{ and } \mathcal{O}_C(C) \text{ has positive degree}) \\
&\geq 2
\end{aligned}$$

So taking 2 sections,  $C$  and one other, we get a rational map  $S \dashrightarrow \mathbb{P}^1$ . After blowing up, this becomes a morphism  $\tilde{S} \dashrightarrow \mathbb{P}^1$ . One of its fibers is isomorphic to  $C$ . By the Noether-Enriques theorem, it follows that  $S$  is rational.  $\square$

So now we want to prove Castelnuovo'. Instead we'll prove

**Castelnuovo''.**  $q = P_2 = 0$  implies that there is an effective divisor  $E$  on  $S$  such that  $K \cdot E < 0$  and  $|K + E| = \emptyset$ . *Keep on board: We seek  $|E| \neq \emptyset$ ,  $|E + K| = \emptyset$ ,  $K \cdot E < 0$ .*

**Castelnuovo'' implies Castelnuovo'.** For then some component  $C$  of  $E$  satisfies  $K \cdot C < 0$ , and any component satisfies  $h^0(S, K + C) = 0$ . Applying Riemann-Roch to  $K + C$  we get

$$\begin{aligned}
0 &= h^0(K + C) \\
&\geq h^0(K + C) - h^1(K + C) + h^0(-C) \\
&= \chi(K + C) \\
&= \chi(\mathcal{O}_X) + \frac{1}{2}((K + C) - K) \cdot (K + C) \\
&> h^0(\mathcal{O}_X) - h^1(\mathcal{O}_X) + h^2(\mathcal{O}_X) + \frac{1}{2}(C + K) \cdot C \\
&\geq 1 + \frac{1}{2}(C + K) \cdot C \\
&= g(C).
\end{aligned}$$

Hence  $g(C) = 0$ .  $(C + K) \cdot C = -2$ , hence  $C^2 \geq -1$ . If  $C^2 = -1$ , then  $C$  is an exceptional curve, and we hypothesized that there weren't any. So Castelnuovo' follows.  $\square$

**Proof of Castelnuovo'' in the case  $K^2 = 0$ .**

How can we possibly use  $P_2 = 0$ ? Only one reasonable way: Our hypothesis  $P_2 = 0$  gives  $h^2(-K) = 0$  (Serre duality). Hence by Riemann-Roch (and  $q = 0$ ):

$$h^0(-K) \geq h^0(-K) - h^1(-K) + h^2(-K) = h^0(\mathcal{O}) - h^1(\mathcal{O}) + h^2(\mathcal{O}) + K^2 \geq 1 + K^2.$$

(We'll use this in the  $K^2 > 0$  case too.)

So  $| -K | \neq \emptyset$ . Let  $H$  be a hyperplane section of  $S$ . Then  $H \cdot K < 0$ . Note:

- If  $n = 0$ , then  $|H + nK| \neq \emptyset$ .
- If  $n \gg 0$  then  $|H + nK| = \emptyset$  (as  $(H + nK) \cdot H < 0$ )

Thus there is an  $n \geq 0$  such that  $|H + nK| \neq \emptyset$ , but  $|H + (n+1)K| = \emptyset$  as  $|H| \neq \emptyset$ , and  $(H + nK) \cdot H < 0$  for  $n \gg 0$ . Let  $D$  be an element.  $|K + D| = \emptyset$ , and  $K \cdot D = -(-K) \cdot H < 0$ .  $\square$

### Proof of Castelnuovo'' in the case $K^2 > 0$ .

Recall  $h^0(-K) = 1 + K^2$ , so  $h^0(-K) \geq -2$ . Suppose  $D \in |-K|$ .

Three cases:

- (1) There is a reducible choice of  $D$ , i.e.  $A, B$  effective with  $A + B \in |-K|$ .
- (2)  $\text{Pic}(C) = \mathbb{Z}K$ . (This implies that there is no reducible choice of  $D$  (why?), but we don't care.)
- (3) All divisors in  $|-K|$  irreducible, and  $\text{Pic}(C) \neq \mathbb{Z}K$ .

*Case 1: There is a reducible choice of  $D$ , i.e.  $A, B$  effective with  $A + B \in |-K|$ . Then  $A \cdot K$  or  $B \cdot K < 0$ , say the former. Then  $A$  is an effective divisor on  $S$  such that  $A \cdot K < 0$ , and  $|A + K| = |-B| = \emptyset$ .*

*Case 2:  $\text{Pic}(C) = \mathbb{Z}K$ . This is the only case where characteristic 0 comes up! From the exact sequence*

$$H^1(S, \mathcal{O}_S) \rightarrow \text{Pic } S \rightarrow H^2(S, \mathbb{Z}) \rightarrow H^2(S, \mathcal{O}_S)$$

*we have  $H^2(S, \mathbb{Z}) \cong \text{Pic } S = \mathbb{Z}K$ . Thus  $b_2 = 1$ . By Poincare duality, the intersection form on  $H^2(S, \mathbb{Z})$  is unimodular, so  $K^2 = 1$ . By Noether's formula,*

$$1 = \chi(\mathcal{O}_S) = \frac{1}{12}(K^2 + 2 - 2b_1 + b_2)$$

*from which  $b_1 = -4$ , contradiction.*

*Case 3: All divisors  $D$  in  $|-K|$  irreducible and  $\text{Pic}(C) \neq \mathbb{Z}K$ . Suppose  $H$  were an effective divisor. As  $|-K| \neq \emptyset$ , there exists  $n > 0$  such that  $|H + nK| \neq \emptyset$  and  $|H + (n+1)K| = \emptyset$ . If  $(H + nK) \cdot K < 0$ , we'd be done.*

*Take an  $H$  such that  $H + nK \neq \emptyset$ . Let  $E \in |H + nK|$ ,  $E = \sum n_i C_i$ . Then  $K \cdot E = -D \cdot E$ , and by the useful remark  $D \cdot E \geq 0$  since  $D$  is irreducible. We are painfully close to being done: we have  $K \cdot E \leq 0$ , and we want  $K \cdot E < 0$ !*

*Thus  $K \cdot C_i \leq 0$  for some  $C = C_i$ . Hence  $|K + C| = \emptyset$ , from which  $0 = h^0(K + C) \geq 1 + \frac{1}{2}(C^2 + CK) = g(C)$ .  $g(C) = 0$ , and  $C^2 = -2 - K \cdot C$  (genus formula). We have gained exactly one thing in this paragraph: our divisor  $C$  is irreducible, whereas our divisor  $E$  was not necessarily. We know that  $|C| \neq \emptyset$ ,  $|K + C| = \emptyset$ , and  $K \cdot C \leq 0$ , and we want to show that  $K \cdot C < 0$ .*

*So we'll assume  $K \cdot C = 0$ , and find a contradiction. From the genus formula,  $C^2 = -2$ . We'll calculate  $h^0(-K - C)$ . Note that  $h^0(2K + C) = h^0(2K + (-D)) \leq h^0(K + C) = 0$ .*

Thus

$$\begin{aligned}
h^0(-K - C) \geq \chi(-K - C) &= \chi(\mathcal{O}_X) + \frac{1}{2}((K + C)^2 + K(K + C)) \\
&= 1 + \frac{1}{2}(C^2 + 3KC + 2K^2) \\
&\geq K^2 \\
&\geq 1
\end{aligned}$$

Since  $C^2 = -2$ , we have  $C \neq -K$ , so there exists a nonzero effective divisor  $A$  such that  $A + C \in |-K|$ . This contradicts our hypothesis that  $|-K|$  has no reducible divisors.

All that's left is:

**Proof of Castelnuovo'' in the case  $K^2 < 0$ .**