

# COMPLEX ALGEBRAIC SURFACES CLASS 17

RAVI VAKIL

## CONTENTS

1. Castelnuovo's Theorem	1
2. Application: The only minimal rational surfaces are $\mathbb{P}^2$ and $\mathbb{F}_n$ ( $n \neq 1$ )	3
3. Albanese variety	4
3.1. The universal property of the Albanese variety	4

On board beforehand:

- Useful trick.  $|D| \neq \emptyset$  (i.e.  $h^0(D) > 0$ ),  $C$  irreducible,  $C^2 \geq 0$  implies  $DC \geq 0$ .
- Genus formula.  $2g(C) - 2 = C \cdot (K_S + C)$ .
- Riemann-Roch:  $\chi(D) = \chi(\mathcal{O}) + \frac{1}{2}D \cdot (D - K)$ .
- Riemann-Roch: In the case when  $h^1(\mathcal{O}) = 0$  and  $h^0(K - D) = 0$ , we have  $h^0(D) \geq 1 + \frac{1}{2}D \cdot (D - K)$  (with equality iff  $h^1(D) = h^2(\mathcal{O}) = 0$ ).

Also, I have been repeatedly using a fact that I may not have explicitly mentioned. A singular curve  $C$  still has a notion of genus (called the *arithmetic genus*),  $g(C) = 1 - \chi(\mathcal{O}_C)$ , which is  $h^1(C, \mathcal{O}_C)$  if  $C$  is connected. (Warning: this is usually denoted  $p_a$ .) Fact:  $C$  irreducible and  $g(C) = 0$  implies that  $C$  is smooth and genus 0.

## 1. CASTELNUOVO'S THEOREM

We spent last time on most of the proof of :

**Theorem: Castelnuovo's Rationality Criterion.** Let  $S$  be a surface with  $q = P_2 = 0$ . Then  $S$  is rational.

*Reminder.*  $q = h^1(S, \mathcal{O}_S) = h^0(S, \Omega_S) = h^2(S, \Omega_S) = h^1(S, K_S)$  (draw Hodge diamond). This is called the *irregularity* of a surface.  $P_2 = h^0(S, K_S^{\otimes 2})$ .

We first reduced this to:

**Castelnuovo'.** Let  $S$  be a minimal surface with  $q = P_2 = 0$ . Then there exists a smooth rational curve  $C$  on  $S$  such that  $C^2 \geq 0$ .

---

*Date:* Wednesday, November 27, 2002. *Patch on Fri. Dec. 5.*

And then reduced it to:

**Castelnuovo".**  $q = P_2 = 0$  and  $S$  minimal implies that there is an effective divisor  $E$  on  $S$  (i.e.  $|E| \neq \emptyset$ ) such that  $K \cdot E < 0$  and  $|K + E| = \emptyset$ . *Keep on board.*

The proof was in 3 cases:  $K^2 < 0$ ,  $= 0$ , and  $> 0$ .

We did the cases  $K^2 = 0$  and  $K^2 > 0$ , so we're left with:

*Proof of Castelnuovo" in the case  $K^2 < 0$ .*

*Step 1. Find an effective divisor  $E$  on  $S$  such that  $K \cdot E < 0$ . (2 out of the 3 desired conclusions.)* Let  $H$  be a hyperplane section of  $S$ . If  $K \cdot H < 0$ , we can take  $E = H$ , and we win. If  $K \cdot H = 0$ , then  $K + nH$  has sections for  $n \gg 0$ , and we win again: take  $E \in |K + nH|$ ; it is effective, and  $K \cdot E < 0$ .

So now assume  $K \cdot H > 0$ . Let  $r_0 = K \cdot H / (-K^2)$ . This is positive. Note that

$$(H + r_0 K)^2 = H^2 + 2(K \cdot H / (-K^2))HK + (K \cdot H)^2 / (-K^2)^2 K^2 = H^2 + (K \cdot H)^2 / (-K^2) > 0.$$

(Don't be nervous with rational coefficients; we can still formally manipulate them in the same way. We should just be careful about them being actual divisors.) Also  $(H + r_0 K) \cdot K = 0$ . So if  $p/q$  ( $p, q \in \mathbb{Z}^+$ ) is a touch greater than  $r_0$ , we have

$$(H + \frac{p}{q}K)^2 > 0, \quad (H + \frac{p}{q}K) \cdot K < 0, \quad (H + \frac{p}{q}K) \cdot H > 0.$$

Let  $D_m = mq(H + \frac{p}{q}K)$ , so

$$D_m^2 > 0, \quad D_m \cdot K < 0, \quad D_m \cdot H > 0.$$

As  $m \rightarrow \infty$ ,  $h^0(K - D_m) = 0$ , as  $(K - D_m) \cdot H < 0$ . By Riemann-Roch,

$$h^0(D_m) \geq \chi(\mathcal{O}_X) + \frac{1}{2}(D_m^2 - D_m \cdot K) \rightarrow \infty$$

as  $m \rightarrow \infty$ . Hence  $h^0(D_m) > 0$  for some large  $m$ ; take  $E \in |D_m|$ . It is effective, and  $K \cdot E < 0$ .

*Step 1. Remember our useful remark from a few classes ago: If  $D$  is an effective divisor, and  $C$  is an irreducible curve such that  $C^2 \geq 0$ , then  $D \cdot C \geq 0$ .*

Suppose we have an effective divisor  $E$  on  $S$  such that  $K \cdot E < 0$ . Then let  $C$  be some irreducible component of  $E$  such that  $K \cdot C < 0$ . By the genus formula,

$$2g(C) - 2 = K \cdot C + C^2 \Rightarrow -2 = K \cdot C + C^2 - 2g(C) \leq -1 + C^2,$$

so  $C^2 \geq -1$ , and  $C^2 = -1$  iff  $C$  is an exceptional curve, which is excluded. Hence  $C^2 \geq 0$ .

Thus  $(aC + nK) \cdot C < 0$  for  $n \gg 0$ . Thus by the useful remark,  $h^0(S, aC + nK) = 0$  for  $n \gg 0$ . Clearly  $h^0(S, aC + 0K) > 0$  (as there is a divisor in the linear system  $aC$ , notably  $aC$  itself), so there is some  $n$  such that  $|aC + nK| \neq \emptyset$ ,  $|aC + (n+1)K| = \emptyset$ . If  $D \in |aC + nK|$ , then  $K \cdot D \leq -a$ , and  $|K + D| = \emptyset$ .  $\square$

## 2. APPLICATION: THE ONLY MINIMAL RATIONAL SURFACES ARE $\mathbb{P}^2$ AND $\mathbb{F}_n$ ( $n \neq 1$ )

Suppose  $S$  is a minimal rational surface. Let  $H$  be a hyperplane section, i.e. a very ample divisor. Consider the set  $A$  of smooth rational curves  $C$  with  $C^2 \geq 0$ . By Castelnuovo', this set is non-empty.

**Castelnuovo'.** Let  $S$  be a minimal surface with  $q = P_2 = 0$ . Then there exists a smooth rational curve  $C$  on  $S$  such that  $C^2 \geq 0$ .

*Keep on board.* Let  $m$  be the smallest possible choice of  $C^2$  in this family. Among those with this value of  $C^2$ , choose some  $C$  with the smallest possible value of  $C \cdot H$  (which is necessarily positive).

**Lemma.** All curves in  $|C|$  are smooth and genus 0.

*Proof.* Suppose  $D = \sum n_i C_i \in |C|$  isn't irreducible.  $K \cdot D < 0$ , so choose some  $C_j$  such that  $K \cdot C_j < 0$ . I'll show you that  $C_j$  is a smooth rational curve with  $C_j^2 \leq C^2$  and  $H \cdot C_j < H \cdot C$ , contradicting our choice of  $C$ .

*Andre pointed out a gap here in the original notes. Here is the patch. By the useful trick, if  $h^0(S, K_S + C) > 0$ , then  $(K_S + C) \cdot C \geq 0$ . But by the genus formula,  $(K_S + C) \cdot C = -2$ .*

Recall:

**Exercise.** (I proved this when showing that Castelnuovo'' implies Castelnuovo', but it's a good exercise to re-prove it yourself.) If  $K_S \cdot C < 0$  and  $h^0(S, K_S + C) = 0$  and  $q = 0$ , then  $C$  is a smooth rational curve with  $C^2 \geq -1$ . Method: apply Riemann-Roch to  $K_S + C$ .

Now here  $q = 0$  (as  $S$  is rational), and  $h^0(S, \mathcal{O}_S(K + C_j)) \leq h^0(S, \mathcal{O}_S(K + \sum n_i C_i)) = 0$ . So  $C_j$  is a smooth rational curve. Also,  $C^2 = (\sum n_i C_i)^2 \geq \sum n_i^2 C_i^2 \geq C_j^2$ . Finally,  $C \cdot H = (\sum n_i C_i) \cdot H \geq C_j \cdot H$ .  $\square$

**Lemma.**  $\dim |C| \leq 2$ , i.e.  $h^0(S, \mathcal{O}(C)) \leq 3$ .

*Proof.* Assume otherwise. Pick some point  $p \in S$ . Pick an open subset containing  $p$ , over which  $\mathcal{O}(C)$  is trivialized. Then via this trivialization, the  $h^0(S, \mathcal{O}(C))$  corresponds to some vector space functions, of dimension at least 4. Thus there is some non-zero function that restricts to something vanishing to order 2 at  $p$ . In that case, the zero set is singular, contradicting our previous lemma.  $\square$

**Lemma.**  $m = 0$  or  $1$ .

*Proof.* We know  $m \geq 0$  by construction. Let  $C_0$  be a fixed curve in that class. Consider the exact sequence  $0 \rightarrow \mathcal{O}_S \rightarrow \mathcal{O}_S(C) \rightarrow \mathcal{O}_{C_0}(m) \rightarrow 0$ . As  $q = 0$ , this is exact on global sections, so  $3 \geq h^0(S, \mathcal{O}_S(C)) = 1 + h^0(\mathbb{P}^1, \mathcal{O}(m)) = 2 + m$ .  $\square$

Note also that  $\mathcal{O}(C)$  has no base points. (Explain.)

**Theorem.** Let  $S$  be a minimal rational surface. Then  $S \cong \mathbb{P}^2$  or some  $\mathbb{F}_n$  for  $n \neq 1$ .

*Proof.* If  $m = 0$ , then  $|C|$  determines a morphism  $S \rightarrow \mathbb{P}^1$  (as there are no base points), all of whose fibres are smooth rational curves. So  $S$  is a rational ruled surface  $\mathbb{F}_n$ .

If  $m = 1$ , then  $|C|$  determines a morphism  $S \rightarrow \mathbb{P}^2$ . The preimages are reduced points (any two sections intersect in  $m = 1$  point, counted with multiplicity), so this is an isomorphism.  $\square$

### 3. ALBANESE VARIETY

I next want to introduce the last important tool you haven't yet seen in the classification of surfaces, the *albanese variety*.

As a goal, I will prove (next day):

**Theorem.** Let  $S, S'$  be two non-ruled minimal surfaces. Then every birational map from  $S'$  to  $S$  is an isomorphism. In particular, every non-ruled surface admits a unique minimal model (up to isomorphism); the group of birational maps from a non-ruled minimal surface to itself coincides with the group of automorphisms of the surface.

**3.1. The universal property of the Albanese variety.** An *abelian variety* over  $\mathbb{C}$  is a projective algebraic variety that is a complex torus  $\mathbb{C}^n/L$  ( $L$  a lattice of rank  $2n$ ); it is a group. In the purely algebraic category, an abelian variety over a field  $k$  is a smooth projective group scheme.

**Hard theorem.** For any (smooth projective) variety  $X$  over a field  $k$ , there exists an abelian variety  $A$  and a morphism  $X \rightarrow A$  with the following universal property: *for any abelian variety  $T$  and any morphism  $f : X \rightarrow T$ , there exists a unique morphism  $\tilde{f} : A \rightarrow T$  such that  $\tilde{f} \circ \alpha = f$ .*

**Exercise.**  $A$  is determined up to isomorphism.

This is called the *Albanese variety* of  $X$ , written  $\text{Alb}(X)$ .

**Hard theorem continued.**  $\alpha$  induces an isomorphism  $\alpha^* : H^0(A, \Omega_A^1) \rightarrow H^0(X, \Omega_X^1)$ .

The proof in general is hard, but over  $\mathbb{C}$  there is an easier proof using Hodge theory (see Beauville, for example). For simplicity, I'll work in the complex category.

**Some fast consequences.**

$\dim \text{Alb}(X) = h^1(X, \Omega_X^1)$ . In particular, if  $X$  is a surface,  $\dim \text{Alb}(X) = q$ . So as a corollary, if  $q = 0$ , every morphism from  $X$  to a complex torus is trivial (i.e. every map is

reduced to a point). Also, every map from  $\mathbb{P}^1$  to a complex torus is trivial. (Explain why in the complex category.)

The Albanese variety is functorial in nature. If  $f : X \rightarrow Y$  is a morphism of smooth projective varieties, there is a unique morphism  $F : \text{Alb}(X) \rightarrow \text{Alb}(Y)$  such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \alpha_X \downarrow & & \downarrow \alpha_Y \\ \text{Alb}(X) & \xrightarrow{\text{exists unique } F} & \text{Alb}(Y) \end{array}$$

is commutative.

Moreover, the Abelian variety  $\text{Alb}(X)$  is generated by  $\alpha(X)$  (i.e. there is no abelian subvariety of  $\text{Alb}(X)$  containing  $\alpha(X)$ ), as the Abelian subvariety of  $\text{Alb}(X)$  generated by  $\alpha(X)$  satisfies the universal property. In particular,  $\alpha(X)$  is not reduced to a point if  $\text{Alb}(X)$  is not a point. It also follows that if the morphism  $f$  is surjective, then so is the morphism  $F : \text{Alb}(X) \rightarrow \text{Alb}(Y)$ .

*Fact:* If  $X$  is a curve, then  $\text{Alb}(X)$  is the Jacobian  $\text{Pic}^0(X)$  (usually denoted  $\text{Jac}(X)$ ).