

# COMPLEX ALGEBRAIC SURFACES CLASS 4

RAVI VAKIL

## CONTENTS

1. Serre duality and Riemann-Roch; back to curves	2
2. Applications of Riemann-Roch	2
2.1. Classification of genus 2 curves	3
2.2. A numerical criterion for very ampleness.	3
3. Normal bundles; the adjunction formula	4
3.1. Applications of the adjunction formula	5
3.2. Classification of curves of genus 3	5

**Recap of last time.** Extended example of  $\mathbb{P}^n$ .  $\mathcal{O}_{\mathbb{P}^n}(d) \leftrightarrow$  degree  $d$  homogeneous things.

Maps to projective space corresponded to vector spaces of sections of an invertible sheaf  $\mathcal{L}$  that are *basepoint free* (no common zero). Hyperplane sections correspond to  $H^0(X, \mathcal{L})$ .

For example, if the sections of  $\mathcal{L}$  have no common zero, then we can map to some projective space by the vector space of *all* sections. Then we say that *the invertible sheaf is basepoint free*. (I didn't give this definition last time.)

**Definition.** The corresponding map to projective space is called a *linear system*. (I'm not sure if I'll use this terminology, but I want to play it safe.)

$$|\mathcal{L}| : X \rightarrow \mathbb{P}^n = \mathbb{P}H^0(X, \mathcal{L})^*.$$

**Definition.** An invertible sheaf  $\mathcal{L}$  is *very ample* if the global sections of  $\mathcal{L}$  gives a closed immersion into projective space.

*Fact.* equivalent to: “separates points and tangent vectors”.

*Definition.* An invertible sheaf is *ample* if some power of it is very ample.

*Note:* A very ample sheaf on a curve has positive degree. Hence an ample sheaf on a curve has positive degree. We'll see later today that this is an “if and only if”.

**Fact (Serre vanishing).** Suppose  $\mathcal{M}$  is any coherent sheaf e.g. an invertible sheaf, or more generally a locally free sheaf (essentially, a vector bundle), and  $\mathcal{L}$  is *ample*. Then for  $n \gg 0$ ,  $H^i(X, \mathcal{M} \otimes \mathcal{L}^n) = 0$  for  $i > 0$ .

## 1. SERRE DUALITY AND RIEMANN-ROCH; BACK TO CURVES

**Fact: Serre duality.** If  $X$  is proper nonsingular and dimension  $n$ , then for  $0 \leq i \leq n$ ,

$$H^i(X, \mathcal{L}) \otimes H^{n-i}(X, \mathcal{K} \otimes \mathcal{L}^\vee) \rightarrow H^n(X, \mathcal{K}) \sim \mathbb{C}$$

is a perfect pairing.

(True for vector bundles. More general formulation for arbitrary coherent sheaves.) Thus  $h^i(X, \mathcal{L}) = h^{n-i}(X, \mathcal{K} \otimes \mathcal{L}^*)$  and  $\chi(C, \mathcal{L}) = (-1)^n \chi(C, \mathcal{K} \otimes \mathcal{L}^\vee)$ .

In particular, we have **Serre duality for curves**. For any invertible sheaf  $\mathcal{L}$ , the map

$$H^0(C, \mathcal{L}) \otimes H^1(C, \mathcal{K} \otimes \mathcal{L}^\vee) \rightarrow H^1(C, \mathcal{K}) \sim \mathbb{C}$$

is a perfect pairing.

Hence two possible definitions of genus are the same:  $h^0(C, \mathcal{K})$  and  $h^1(C, \mathcal{O}_C)$ .

**Fact: Riemann-Roch Theorem.**

$$h^0(C, \mathcal{L}) - h^1(C, \mathcal{L}) = \deg(\mathcal{L}) - g + 1.$$

(I.e.  $\chi(C, \mathcal{L}) = \deg - g + 1$  — remember that the cohomology of a coherent sheaf vanishes above the dimension of the variety.)

Generalizations: Hirzebruch-Riemann-Roch, to Grothendieck-Riemann-Roch. Hirzebruch-Riemann-Roch, which we'll be using, is a consequence of the Atiyah-Singer index theorem, which Rafe Mazzeo spoke about in the colloquium yesterday.

Proof of Riemann-Roch: (i) algebraic, (ii) complex-analytic, (iii) Atiyah-Singer.

Using Serre duality:

$$h^0(C, \mathcal{L}) - h^0(C, \mathcal{K} \otimes \mathcal{L}^\vee) = \deg(\mathcal{L}) - g + 1.$$

**Corollary.**  $\deg K = 2g - 2$ . (Do it.)

So we have another definition of  $g$ .

## 2. APPLICATIONS OF RIEMANN-ROCH

**Theorem.** If  $\deg \mathcal{L} \geq 2g - 1$ , then  $h^1(C, \mathcal{L}) = 0$ . Hence  $h^0(C, \mathcal{L}) = \deg \mathcal{L} - g + 1$ .

Proof: Serre duality, and the fact that invertible sheaves of negative degree have no sections. Then Riemann-Roch.

**Theorem (numerical criterion for basepoint freeness).** If  $\deg \mathcal{L} \geq 2g$ , then  $\mathcal{L}$  is basepoint free. (Remind them of definition.)

*Proof:*  $H^0(C, \mathcal{L}(-p))$  is the vector space of sections with a zero at  $p$ . Want to show that  $H^0(C, \mathcal{L}) - H^0(C, \mathcal{L}(-p)) > 0$ . Do it using previous result: difference is 1!

**Example.** If  $C$  is genus 1 and  $\mathcal{L}$  is degree 2, then we get a map to  $\mathbb{P}^1$ . This shows that every genus 1 curve is a double cover of  $\mathbb{P}^1$ . (Explain “hyperplane section” in this case.)

**2.1. Classification of genus 2 curves. Theorem.** Any genus 2 curve has a unique double cover of  $\mathbb{P}^1$  branched over 6 points. (The 6 points are unique up to automorphisms of  $\mathbb{P}^1$ .) Any double cover of  $\mathbb{P}^1$  branched over 6 points comes from a genus 2 curve. Hence genus 2 curves are classified by the space of 6 distinct points of  $\mathbb{P}^1$ , modulo automorphisms of  $\mathbb{P}^1$ . In particular, there is a dimension  $6 - \dim \text{Aut } \mathbb{P}^1 = 3$  moduli space.

*Proof.* First, every genus 2 curve  $C$  has a degree 2 map to  $\mathbb{P}^1$  via  $\mathcal{K}$  (basepoint freeness). The numerical criterion for basepoint freeness doesn’t apply, unfortunately. Let’s check that  $\mathcal{K}$  has 2 sections:  $h^0(\mathcal{K}) - h^0(\mathcal{O}) = \deg(\mathcal{K}) - 2 + 1 = 1$ . Next, let’s check that  $\mathcal{K}$  is basepoint free. We want  $\mathcal{K}(-p)$  has only 1 section for all  $p$ . If  $\mathcal{K}(-p)$  had two sections, then we’d have a degree 1 map to  $\mathbb{P}^1$  from  $C$  (explain), contradiction.

Next we’ll see that the only degree 2 maps to  $\mathbb{P}^1$  arise from the canonical bundle. Suppose  $\mathcal{L}$  is degree 2, and has 2 sections. Then by RR,  $\mathcal{K} \otimes \mathcal{L}^*$  has 1 section, and is degree 0. But the only degree 0 sheaf with a section is the trivial sheaf.

Finally, any double cover of  $\mathbb{P}^1$  branched over 6 points is genus 2. This follows from the *Riemann-Hurwitz formula*.

Using naive geometry:  $2\chi(\mathbb{P}^1) - 6 = \chi(C)$ , i.e.  $-2 = 2 - 2g$ , so  $g = 2$ .

*Fact.* Riemann-Hurwitz formula:  $\mathcal{K}_C = \pi^* \mathcal{K}_{\mathbb{P}^1} + \text{ramification divisor}$ . Taking degrees:  $2g - 2 = 2(-2) + 6$ . □

**2.2. A numerical criterion for very ampleness. Theorem.** If  $\deg \mathcal{L} \geq 2g + 1$ , then  $\mathcal{L}$  is very ample. (Remind them of definition.)

*Proof.* I’ll show you that it separates points. Suppose  $p \neq q$ .  $H^0(C, \mathcal{L}) - H^0(C, \mathcal{L}(-p - q)) = 2$ . Hence there is a section vanishing at  $p$  and not at  $q$ , and vice versa.

Same idea works for separating tangent vectors.  $h^0(C, \mathcal{L}) - h^0(C, \mathcal{L}(-2p)) = 2$ . Hence there is a section vanishing at  $p$  but not to order 2.

**Example.**  $C$  genus 1,  $\mathcal{L}$  has degree 3. Then  $C \hookrightarrow \mathbb{P}^2$ , degree 3. Hence every elliptic curve can be described by a plane cubic. (We’ll soon see that any plane cubic is a genus 1 curve.)

**Corollary.**  $\mathcal{L}$  on  $C$  is ample iff it has positive degree.

### 3. NORMAL BUNDLES; THE ADJUNCTION FORMULA

There is one more natural place to find line bundles: the normal bundle to a submanifold of dimension 1 less. I'll now discuss normal bundles algebraically, and as an application prove the *adjunction formula*.

Suppose  $D$  is a nonsingular divisor (codimension 1) on nonsingular  $X$ . Then we can understand the canonical sheaf of  $D$  in terms of the canonical sheaf of  $X$ .

**The adjunction formula.**  $\mathcal{K}_D = \mathcal{K}_X(D)|_D$ .

(Remind them what  $\mathcal{K}_X(D)$  means.)

This actually holds in much more generality, e.g.  $D$  can be arbitrarily singular, and  $X$  need not be smooth.

Here is an informal description of why this is true.

- Motivation: tangent space in differential geometry to a point  $p$  in a manifold  $W$  is the space of curves, modulo some equivalence relation.

- The cotangent space is the dual of this space, and can be interpreted as the space of functions vanishing at  $p$  modulo an equivalence relation: the functions vanishing to order 2 at  $p$ . In the space  $\mathcal{O}$  of functions near  $p$ , the first is the ideal  $I_p$ , the second is the ideal  $I_p^2$ . Thus the cotangent space is  $I_p/I_p^2$ .

- Aside: in this case,  $I_p$  is a maximal ideal  $\mathfrak{m}_p$ , as  $\mathcal{O}/I_p$  is a field. So the cotangent space is  $\mathfrak{m}_p/\mathfrak{m}_p^2$ . That's why the Zariski tangent space is defined as  $\mathfrak{m}_p/\mathfrak{m}_p^2$ . This is a purely algebraic definition. It works for any algebraic  $X$ , not even defined over a field, not even non-singular.

- Next consider  $Y$  to be a nonsingular subvariety of  $X$ , of codimension  $c$ . Then there is a conormal bundle of  $Y$  in  $X$  (the dual of the normal bundle. It has rank  $d$ . The conormal sheaf is the sheaf of sections of  $Y$ . It is locally free of rank  $d$ . *Definition (motivated by previous discussion)*. In sheaf language: the conormal bundle is  $\mathcal{I}/\mathcal{I}^2$ . This is a priori a sheaf on all of  $X$ , but in fact it lives on  $Y$  ("is supported on  $Y$ ").

- Now suppose further that  $Y$  is a divisor, so I'll now call it  $D$ . Then  $\mathcal{I} = \mathcal{O}_X(D)$ . We're modding out this sheaf by functions vanishing on  $D$ ; this is the same as restricting to  $D$ . Hence  $\mathcal{I}/\mathcal{I}^2 \sim \mathcal{O}(D)|_D$ .

*Proof of the adjunction formula.* We have an exact sequence

$$0 \rightarrow T_D \rightarrow T_X|_D \rightarrow N_D \rightarrow 0.$$

Dualize to get:

$$0 \rightarrow N_D^* \rightarrow \Omega_X|_D \rightarrow \Omega_D \rightarrow 0.$$

Take top wedge powers to get  $\mathcal{K}_X|_D \sim \mathcal{K}_D \otimes N_D^*$  from which

$$\mathcal{K}_D \sim \mathcal{K}_X|_D \otimes N_D = \mathcal{K}_X|_D \otimes \mathcal{O}_X(D)|_D = \mathcal{K}_X(D)|_D.$$

□

**3.1. Applications of the adjunction formula.** 1) Cubics in  $\mathbb{P}^2$  have trivial canonical bundle. Hence all genus 1 curves have canonical sheaf that is not degree 0, but also trivial.

2) Quartics in  $\mathbb{P}^3$  also have trivial canonical bundle. K3 surfaces.

3) What's the genus of a smooth degree  $d$  curve in  $\mathbb{P}^3$ ? Answer:  $(d - 1)(d - 2)/2$ .

4) Smooth quartics in  $\mathbb{P}^2$  are embedded by their canonical sheaf. More on this in a second.

5) Smooth complete intersection of surfaces of degree 2 and 3 in  $\mathbb{P}^3$ : also embedded by their canonical sheaf.

**3.2. Classification of curves of genus 3.** I'll discuss at greater length next day.

**Theorem.** Every smooth curve of genus 3 is of (precisely) one of the two following forms.

(i) A smooth quartic curve in  $\mathbb{P}^2$  (in a unique way, up to automorphisms of  $\mathbb{P}^2$ ).

(ii) A double cover of  $\mathbb{P}^1$  branched over 8 points (in a unique way, up to automorphisms of  $\mathbb{P}^2$ ).